arXiv:1007.1296v2 [math.AG] 29 Nov 2010

AUSLANDER–REITEN DUALITY AND MAXIMAL MODIFICATIONS FOR NON-ISOLATED SINGULARITIES.

OSAMU IYAMA AND MICHAEL WEMYSS

Abstract. We first generalize classical Auslander–Reiten duality for isolated singularities to cover singularities with a one-dimensional singular locus. We then define the notion of CT modules for non-isolated singularities and we show that these are intimately related to noncommutative crepant resolutions (NCCRs). When R has isolated singularities, CT modules recover the classical notion of cluster tilting modules but in general the two concepts differ. Then, wanting to generalize the notion of NCCRs to cover partial resolutions of Spec R, in the main body of this paper we introduce a theory of modifying and maximal modifying modules for three-dimensional Gorenstein rings R. Under mild assumptions all the corresponding endomorphism algebras of the maximal modifying modules are shown to be derived equivalent. We then develop a theory of mutation for modifying modules which is similar but different to mutations arising in cluster tilting theory. The behavior of our mutation strongly depends on whether a certain factor algebra is artinian — when it is not artinian our mutation may be the identity.

Contents

1. Introduction 1
1.1. Motivation and History 1
1.2. Auslander–Reiten Duality for Non-Isolated Singularities 3
1.3. Maximal Modifications and NCCRs 4
1.4. Mutation of Modifications 6
1.5. Conventions 7
2. Preliminaries 7
2.1. Depth and CM Modules 7
2.2. Reflexive Equivalence and Symmetric Algebras 8
2.3. Non-Singular and Gorenstein Orders 9
2.4. d-CY$^-$ Algebras 11
3. Auslander–Reiten Duality for Non-Isolated Singularities 12
4. Modifying and Maximal Modifying Modules 15
5. Relationship between CT modules, NCCRs and MM modules 20
5.1. CT Modules on Non-Isolated Singularities 20
5.2. Modifying Modules on Isolated Singularities 24
6. Mutations of Modifying Modules 25
6.1. General Definitions and First Properties 25
6.2. Complete Local Case 30
References 32

1. Introduction

1.1. **Motivation and History.** One of the basic results in representation theory of commutative algebras is Auslander–Reiten (=AR) duality for isolated singularities, which gives us many important consequences, e.g. the existence of almost split sequences and the Calabi-Yau property of the stable categories of Cohen–Macaulay (=CM) modules over Gorenstein isolated singularities. One of the aims of this paper is to establish AR duality for singularities with one dimensional singular loci. As an application, the stable

---

The second author was partially supported by a JSPS Postdoctoral Fellowship and by the EPSRC.

categories of CM modules over Gorenstein singularities with one dimensional singular loci enjoy a generalized Calabi-Yau property. This is a starting point of our reseach to apply the methods of cluster tilting in representation theory to study singularities.

One of the highlights of representation theory of commutative algebras is AR theory of simple surface singularities. They have only finitely many indecomposable CM modules, and the Auslander algebras (i.e. the endomorphism algebras of the direct sums of all indecomposable CM modules) enjoy many nice properties. If we consider singularities of dimension greater than two, then there are very few representation-finite singularities, and their Auslander algebras do not satisfy such nice properties. The reason is that the categories of CM modules do not behave nicely in the sense that the homological properties of simple functors corresponding to free modules are very hard to control. Motivated to obtain the correct category on which higher AR theory should be performed, in [Iya07] the first author introduced the notion of a maximal n-orthogonal subcategory and maximal n-orthogonal module for the category $\operatorname{mod}\Lambda$, later renamed cluster tilting subcategories and cluster tilting modules respectively. Just as classical AR theory on $\operatorname{mod}\Lambda$ was moved to AR theory on $\operatorname{CM}\Lambda$ following the work of Auslander on the McKay correspondence for surfaces Λ [Aus86], this suggests that in the case of a higher dimensional CM singularity R we should apply the definition of a maximal n-orthogonal subcategory/modules to $\operatorname{CM} R$ and hope that this provides a suitable framework for tackling higher-dimensional geometric problems. Strong evidence for this is provided when R is a three dimensional normal isolated Gorenstein singularity, since in this case it is known that such objects have an int imate relationship with Van den Bergh's noncommutative crepant resolutions (NCCRs) [IR08, 8.13]. Requiring R to be isolated is absolutely crucial to this relationship (by normality the singularities are automatically isolated in the surfaces case); from an algebraic perspective this perhaps should not be surprising since AR theory only works well for isolated singularities. It turns out that the study of maximal n-orthogonal modules in $\operatorname{CM} R$ is not well-suited to non-isolated singularities since the Ext vanishing condition is far too strong; the question arises as to what subcategories of $\operatorname{CM} R$ should play the role of the maximal n-orthogonal subcategories above.

Although in this paper we answer this question, in fact we say much more. Philosophically, the point is that we are asking ourselves the wrong question. The restriction to studying maximal orthogonal modules is unsatisfactory since crepant resolutions need not exist (even for 3-folds) and so we develop a theory which can deal with singularities in the crepant partial resolutions. Since the endomorphism rings of maximal orthogonal modules have finite global dimension, these will not do the job for us.

We introduce the notion of maximal modifying modules (see 1.11 below) which intuitively we think of as corresponding to shadows of maximal crepant partial resolutions. Geometrically this level always exists, but only sometimes will it be smooth. View regards to this viewpoint maximal modifying modules are a more natural class of objects to work with compared to noncommutative crepant resolutions; we should thus always work in this level of generality and simply view the case when the geometry is smooth as being a happy coincidence. Pushing this philosophy further, everything that we are currently able to do with NCCRs we should be able to do with maximal modifying modules, and this motivates much of the work in this paper.

In fact in many regards restricting our attention to only studying maximal crepant partial resolutions misses much of the picture and so we should (and do) work even more generally. When one wants to flop curves between varieties with canonical singularities which are not terminal this does not take place on the maximal level but we should still be able to understand this homologically. This motivates our definition and the study of modifying modules (see 1.11 below). Viewing our modifying modules M as conjectural shadows of partial resolutions we should thus be able to track the birational transformations between the geometrical spaces by using some kind of homological transformation between the corresponding modifying modules. This leads us to develop a theory of mutation for modifying modules, which we do in Section 6.

For the algebraists, we remark that in moving from isolated to non-isolated singularities we must give up the idea of rigidity and so our theory of mutation loses some of the fundamental features from the world of clusters, such as the unique exchange property.

Certain aspects remain however; for example mutating twice at the same vertex always returns us to the original position and furthermore our mutation induces canonical derived equivalences between the corresponding endomorphism rings. We believe that it is precisely the lack of unique exchange together with the induced derived equivalences that provides the level of freedom which is required if the theory is to match (and in fact imply) what is occurring geometrically.

Although in this paper we are principally interested in the geometrical and commutative algebraic statements, the proofs of our theorems require a slightly more general noncommutative setting. For this reason we deal with noncommutative d-CY$^-$ algebras Λ (see §2.4) as a generalization of commutative Gorenstein d-dimensional rings. By doing this we increase the technical difficulty, but we emphasize that we are unable to prove the purely commutative statements without passing to the noncommutative world.

We now describe our results rigorously, and in more detail.

1.2. **Auslander–Reiten Duality for Non-Isolated Singularities.** Throughout this subsection let R be an equi-codimensional (see 1.2 below) CM ring of dimension d with a canonical module ω_R. Recall that for a non-local CM ring R, a finitely generated R-module ω_R is called a *canonical module* if $(\omega_R)_{\mathfrak{m}}$ is a canonical $R_{\mathfrak{m}}$-module for all $\mathfrak{m} \in \operatorname{Max} R$ [BH, 3.3.16]. In this case $(\omega_R)_{\mathfrak{p}}$ is a canonical $R_{\mathfrak{p}}$-module for all $\mathfrak{p} \in \operatorname{Spec} R$ since canonical modules localize for local CM rings [BH, 3.3.5].

We denote $\operatorname{CM} R$ to be the category of CM R-modules (see §2.1), $\underline{\operatorname{CM}}R$ to be the *stable category* and $\overline{\operatorname{CM}}R$ to be the *costable category*. The *AR translation* is defined to be

$$\tau := \operatorname{Hom}_R(\Omega^d \operatorname{Tr}(-), \omega_R) : \underline{\operatorname{CM}}R \to \overline{\operatorname{CM}}R.$$

When R is an isolated singularity one of the fundamental properties of the category $\operatorname{CM} R$ is the existence of Auslander–Reiten duality [Aus78, 8.8] [AR], namely

$$\underline{\operatorname{Hom}}_R(X, Y) \cong D \operatorname{Ext}^1_R(Y, \tau X)$$

for all $X, Y \in \operatorname{CM} R$ where D is the Matlis dual (see §3). We generalize this to mildly non-isolated singularities by proving the following theorem:

Theorem 1.1. *(=3.1) Let R be a d-dimensional, equi-codimensional CM ring with a canonical module ω_R and singular locus of Krull dimension less than or equal to one. Then there exists a functorial isomorphism*

$$\mathsf{fl}\, \underline{\operatorname{Hom}}_R(X, Y) \cong D(\mathsf{fl}\, \operatorname{Ext}^1_R(Y, \tau X))$$

for all $X, Y \in \operatorname{CM} R$, where for an R-module M we denote $\mathsf{fl}\, M$ to be the largest finite length R-submodule of M.

In fact we prove 3.1 in the setting of noncommutative R-orders (see §3 for precise details). In the above and throughout this paper, for many of the global-local arguments to work we often have to add the following mild technical assumption.

Definition 1.2. *A commutative ring R is* equi-codimensional *if all its maximal ideals have the same height.*

Although technical, such rings are very common; for example all affine domains are equi-codimensional [Ei95, 13.4]. Since our main applications are three-dimensional normal affine domains, in practice this adds no restrictions to what we want to do. We will use the following well-known property [Mat, 17.3(i), 17.4(ii)]:

Lemma 1.3. *Let R be an equi-codimensional CM ring, and let $\mathfrak{p} \in \operatorname{Spec} R$. Then*

$$\operatorname{ht} \mathfrak{p} + \dim R/\mathfrak{p} = \dim R.$$

The above generalized Auslander–Reiten duality implies the following generalized 2-Calabi-Yau property of the triangulated category $\underline{\operatorname{CM}}R$.

Corollary 1.4. *(=3.8) Let R be a normal, 3-dimensional, equi-codimensional Gorenstein ring. Then*

(1) *There is a functorial isomorphism*

$$\mathsf{fl}\, \operatorname{Ext}^1_R(X, Y) \cong D(\mathsf{fl}\, \operatorname{Ext}^1_R(Y, X))$$

for all $X, Y \in \operatorname{CM} R$.
(2) $\operatorname{Hom}_R(X,Y) \in \operatorname{CM} R$ *if and only if* $\operatorname{Hom}_R(Y,X) \in \operatorname{CM} R$.

Note that 1.4(2) also holds for arbitrary dimension (see 2.8). This symmetry in the Hom groups gives us the technical tool we require to move successfully from the cluster tilting level to the maximal modification level below, and is entirely analogous to the symmetry given by [CB, Lemma 1] as used in the work of Geiss–Leclerc–Schröer [GLS].

1.3. Maximal Modifications and NCCRs.

Here we introduce our main definitions, namely modifying, maximal modifying and CT modules, and then survey our main results.

Throughout, an R-algebra is called *module finite* if it is a finitely generated R-module. Recall [Aus78, Aus84, CR90]:

Definition 1.5. *Let R be a CM ring and let Λ be a module finite R-algebra. We say*
(1) *Λ is an* R-order *if* $\Lambda \in \operatorname{CM} R$.
(2) *An R-order Λ is* non-singular *if* $\operatorname{gl.dim} \Lambda_{\mathfrak{p}} = \dim R_{\mathfrak{p}}$ *for all primes $\mathfrak{p}$ of R.*
(3) *An R-order Λ has* isolated singularities *if $\Lambda_{\mathfrak{p}}$ is a non-singular $R_{\mathfrak{p}}$-order for all non-maximal primes $\mathfrak{p}$ of R.*

In the definition of non-singular R-order above, $\operatorname{gl.dim} \Lambda_{\mathfrak{p}} = \dim R_{\mathfrak{p}}$ means that $\operatorname{gl.dim} \Lambda_{\mathfrak{p}}$ takes the smallest possible value. In fact for an R-order Λ we always have that $\operatorname{gl.dim} \Lambda_{\mathfrak{p}} \geq \dim R_{\mathfrak{p}} := t_{\mathfrak{p}}$ for all primes $\mathfrak{p}$ of R since $\operatorname{proj.dim}_{\Lambda_{\mathfrak{p}}} \Lambda_{\mathfrak{p}}/(x_1, ..., x_{t_{\mathfrak{p}}})\Lambda_{\mathfrak{p}} \geq \dim R_{\mathfrak{p}}$ for a $\Lambda_{\mathfrak{p}}$-regular sequence $x_1, \dots, x_{t_{\mathfrak{p}}}$.

Throughout this paper we denote $(-)^* := \operatorname{Hom}_R(-,R) : \operatorname{mod} R \to \operatorname{mod} R$ and we say that $X \in \operatorname{mod} R$ is *reflexive* if the natural map $X \to X^{**}$ is an isomorphism. We denote $\operatorname{ref} R$ to be the category of reflexive R-modules. By Serre, when R is a normal domain the category $\operatorname{ref} R$ is closed under both kernels and extensions.

Definition 1.6. *Let R be CM, then by a* noncommutative crepant resolution *(NCCR) of R we mean $\Gamma := \operatorname{End}_R(M)$ where $M \in \operatorname{ref} R$ is non-zero such that Γ is a non-singular R-order.*

We show in 2.14 that under very mild assumptions the condition in 1.5(2) can in fact be checked at only maximal ideals, and we show in 2.19 that 1.6 is equivalent to the definition of NCCR due to Van den Bergh [V04b] when R is a Gorenstein normal domain. Recall the following:

Definition 1.7. *Let A be a ring. We say that an A-module M is a* generator *if M contains A as a summand.*

Motivated by wanting a characterization of the reflexive generators which give NCCRs, we define:

Definition 1.8. *Let R be a d-dimensional CM ring with a canonical module ω_R. We call $M \in \operatorname{CM} R$ a* CT module *if*

$$\operatorname{add} M = \{X \in \operatorname{CM} R : \operatorname{Hom}_R(M,X) \in \operatorname{CM} R\} = \{X \in \operatorname{CM} R : \operatorname{Hom}_R(X,M) \in \operatorname{CM} R\}.$$

Clearly a CT module is always a generator and cogenerator (i.e. it contains both R and ω_R as summands). We show in 5.16 that this recovers the established notion of maximal 1-orthogonal modules when R is 3-dimensional and has isolated singularities. The following result is shown in [Iya07, 2.5] under the assumption that G is a small subgroup of $\operatorname{GL}(d,k)$ and S^G is an isolated singularity. We can drop both assumptions under our definition of CT modules:

Theorem 1.9. *(=5.9) Let k be a field of characteristic zero and let $S := k[[x_1, \dots, x_d]]$ be the formal power series ring. For an arbitrary finite group G together with a group homomorphism $G \to \operatorname{Aut}_{k\text{−alg}}(S)$, let $R := S^G$. Then S is a CT R-module.*

One of our main results involving CT modules is the following, where part (2) answers a question of Van den Bergh.

Theorem 1.10. *(=5.12) Let R be a 3-dimensional Gorenstein equi-codimensional normal domain. Then*
(1) *CT modules are precisely those reflexive generators which give NCCRs.*

(2) *R has a NCCR* $\iff$ *R has a NCCR given by a CM generator M* $\iff$ $\operatorname{CM} R$ *contains a CT module.*

However R need not have a NCCR so we must weaken the notion of CT module and allow for our endomorphism rings to have infinite global dimension. We do so as follows:

Definition 1.11. *Let R be a d-dimensional CM ring. We call $N \in \operatorname{ref} R$ a* modifying module *if $\operatorname{End}_R(N) \in \operatorname{CM} R$, whereas we call N a* maximal modifying (MM) module *if N is modifying and furthermore it is maximal with respect to this property, that is to say if there exists $X \in \operatorname{ref} R$ with $N \oplus X$ modifying, necessarily $X \in \operatorname{add} N$. Equivalently, we say N is maximal modifying if*

$$\operatorname{add} N = \{X \in \operatorname{ref} R : \operatorname{Hom}_R(N \oplus X, N \oplus X) \in \operatorname{CM} R\}.$$

If N is a MM module, we call $\operatorname{End}_R(N)$ a maximal modification algebra.

When R is 3-dimensional with isolated singularities we show in 5.16 that modifying modules recover the established notion of rigid modules, whereas MM modules recover the notion of maximal rigid modules. However, other than pointing out this relationship (in §5.2), throughout we never assume that R has isolated singularities.

We prove that when R has a NCCR the maximal modifying modules in fact characterize NCCRs:

Proposition 1.12. *(=5.14) Let R be a 3-dimensional Gorenstein equi-codimensional normal domain, and assume that R has a NCCR (equivalently, by 1.10, a CT module). Then*
(1) *MM modules are precisely the reflexive modules which give NCCRs.*
(2) *MM modules which are CM (equivalently, by 4.2, the MM generators) are precisely the CT modules.*
(3) *CT modules are precisely those CM modules which give NCCRs.*

The point is that R need not have a NCCR and our definition of maximal modification algebra is strictly more general. Throughout this paper we freely use the notion of a tilting module which we *always* assume has projective dimension less than or equal to one:

Definition 1.13. *Let Λ be a ring. Then $T \in \operatorname{mod} \Lambda$ is called a* tilting module *if $\operatorname{proj.dim}_\Lambda T \leq 1$, $\operatorname{Ext}^1_\Lambda(T, T) = 0$ and further there exists an exact sequence*

$$0 \to \Lambda \to T_0 \to T_1 \to 0$$

with each $T_i \in \operatorname{add} T$.

Our next result details the relationship between modifying and maximal modifying modules on the level of derived categories.

Theorem 1.14. *(=4.9) Let R be a 3-dimensional Gorenstein equi-codimensional normal domain with MM module M. Then*
(1) *If N is any modifying module then the partial tilting $\operatorname{End}_R(M)$-module $T := \operatorname{Hom}_R(M, N)$ induces a recollement* [BBD]

$$\begin{tikzcd} \mathrm{U} \arrow[r] & \mathrm{D}(\operatorname{Mod} \operatorname{End}_R(M)) \arrow[l, shift right=2] \arrow[l, shift left=2] \arrow[r, "F"] & \mathrm{D}(\operatorname{Mod} \operatorname{End}_R(N)) \arrow[l, shift right=2] \arrow[l, shift left=2] \end{tikzcd}$$

where $F = \mathbf{R}\operatorname{Hom}(T, -)$ and U is a certain triangulated subcategory of $\mathrm{D}(\operatorname{Mod} \operatorname{End}_R(M))$.
(2) *If further N is maximal modifying then the above functor F is an equivalence.*

Recall that two rings A and B are said to be *derived equivalent* if $\mathrm{D}(\operatorname{Mod} \operatorname{End}_R(N)) \approx \mathrm{D}(\operatorname{Mod} \operatorname{End}_R(M))$ [R89]. Theorem 1.14 now immediately gives the following, which we view as the noncommutative analogue of a result of Chen [C02].

Corollary 1.15. *Let R be a 3-dimensional Gorenstein equi-codimensional normal domain with a MM module. Then all maximal modification algebras are derived equivalent.*

In our study of modifying modules, reflexive modules over noncommutative R-algebras play a crucial role.

Definition 1.16. *Let R be any commutative ring. If A is any R-algebra then we say that $M \in \operatorname{mod} A$ is a* reflexive A-module *if it is reflexive as an R-module.*

Note that we do not require that the natural map $M \to \operatorname{Hom}_{A^{\mathrm{op}}}(\operatorname{Hom}_A(M,A),A)$ is an isomorphism however when A is 3-CY$^-$ (see §2.4) and M is a reflexive A-module, automatically it is. Our main theorem regarding maximal modifying modules is the following remarkable relationship between modifying modules and tilting modules. Note that (3) below says that R has a maximal modification algebra if and only if it has a maximal modification algebra $\operatorname{End}_R(N)$ where N is a CM generator, a generalization of 1.10(2).

Theorem 1.17. *(=4.15, 4.16) Let R be a 3-dimensional Gorenstein equi-codimensional normal domain with a maximal modifying module M. Then*
(1) *The functor $\operatorname{Hom}_R(M,-) : \operatorname{mod} R \to \operatorname{mod} \operatorname{End}_R(M)$ induces bijections*

$$\begin{aligned} \{\textit{maximal modifying } R\textit{-modules}\} &\overset{1:1}{\longleftrightarrow} \{\textit{reflexive tilting } \operatorname{End}_R(M)\textit{-modules}\}. \\ \{\textit{modifying } R\textit{-modules}\} &\overset{1:1}{\longleftrightarrow} \{\textit{reflexive partial tilting } \operatorname{End}_R(M)\textit{-modules}\}. \end{aligned}$$

(2) *N is modifying $\iff$ N is the direct summand of a maximal modifying module.*
(3) *R has a MM module containing R as a summand. Necessarily this is CM.*

1.4. **Mutation of Modifications.** Let R be a normal, Gorenstein, three-dimensional, equi-codimensional ring. We introduce categorical mutations as a method of producing modifying modules from a given one. Recall:

Definition 1.18. *For R-modules M and N, we say that a morphism $f : N_0 \to M$ is a* right $\operatorname{add} N$-approximation *if $N_0 \in \operatorname{add} N$ and further*

$$\operatorname{Hom}_R(N, N_0) \xrightarrow{\cdot f} \operatorname{Hom}_R(N, M) \to 0$$

is exact. Dually we define a left $\operatorname{add} N$-approximation.

Now for a given modifying module M, and N such that $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$ we consider

(1) a right $\operatorname{add} N$-approximation of M, denoted $N_0 \xrightarrow{a} M$.
(2) a right $\operatorname{add} N^*$-approximation of M^*, denoted $N_1^* \xrightarrow{b} M^*$.

Note that the above a and b are surjective if N is a generator. In what follows we denote the kernels by

$$0 \to K_0 \xrightarrow{c} N_0 \xrightarrow{a} M \quad \text{and} \quad 0 \to K_1 \xrightarrow{d} N_1^* \xrightarrow{b} M^*$$

and call these *exchange sequences.*

Definition 1.19. *With notation as above, we define the* right mutation *of M at N to be $\mu_N(M) := N \oplus K_0$ and we define the* left mutation *of M at N to be $\nu_N(M) := N \oplus K_1^*$.*

Note that by definition $\nu_N(M) = (\mu_{N^*}(M^*))^*$.

Proposition 1.20. *(=6.2, 6.4)* (1) *Both $\mu_N(M)$ and $\nu_N(M)$ are modifying R-modules.*
(2) *μ and ν are mutually inverse operations, i.e. we have that $\nu_N(\mu_N(M)) = M$ and $\mu_N(\nu_N(M)) = M$, up to additive closure.*

Under fairly weak assumptions it turns out that left mutation is the same as right mutation. If $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$ then we define $[N]$ to be the two-sided ideal of $\operatorname{End}_R(M)$ consisting of morphisms $M \to M$ which factor through a member of $\operatorname{add} N$. Our first main theorem is the following:

Theorem 1.21. *(=6.6, 6.8) Suppose R is a 3-dimensional Gorenstein equi-codimensional normal domain with modifying module M. Suppose $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$. Denote $\Lambda = \operatorname{End}_R(M)$, and define $\Lambda_N := \Lambda/[N]$. If Λ_N is artinian then*
(1) *$\nu_N(M)$ is also a modifying module. Further*
  (a) *If M is maximal modifying then $\nu_N(M)$ is maximal modifying.*
  (b) *If $\operatorname{End}_R(M)$ is a NCCR then $\operatorname{End}_R(\nu_N(M))$ is a NCCR.*
  (c) *If M is a CT module and N is a generator then $\nu_N(M)$ is a CT module.*
(2) *If $\operatorname{add} N \neq \operatorname{add} M$ then $\nu_N(M) \neq M$.*
(3) *$\operatorname{End}_R(M)$ and $\operatorname{End}_R(\nu_N(M))$ are derived equivalent.*
An identical statement holds throughout if we replace $\nu_N(M)$ by $\mu_N(M)$.

Note that when R is finitely generated over a field k, Λ_N is artinian if and only if it is finite dimensional over k (see 6.7). We emphasize that in our proofs, the homological algebra always splits into two cases depending on whether or not Λ_N is artinian. The above 1.21 covers the case when Λ_N is artinian, whereas the case when Λ_N is not artinian is a little more tricky. Nevertheless for maximal modifying modules the picture remains remarkably clear, provided that R is complete local and we mutate at only one indecomposable summand at a time:

Theorem 1.22. *(=6.14) Suppose R is a complete local normal three-dimensional Gorenstein ring with maximal modifying module M. Denote $\Lambda = \operatorname{End}_R(M)$, let M_i be an indecomposable summand of M and consider $\Lambda_i := \Lambda/\Lambda(1-e_i)\Lambda$ where e_i is the idempotent in Λ corresponding to M_i. To ease notation denote $\mu_i = \mu_{\frac{M}{M_i}}$ and $\nu_i = \nu_{\frac{M}{M_i}}$. Then*
(1) If Λ_i is not artinian then $\mu_i(M) = \nu_i(M) = M$.
(2) If Λ_i is artinian then $\mu_i(M) = \nu_i(M)$ and this is not equal to M.
In either case denote $\mu_i := \mu_i = \nu_i$ then it is also true that
(3) $\mu_i\mu_i(M) = M$.
(4) $\mu_i(M)$ is a maximal modifying module.
(5) $\operatorname{End}_R(M)$ and $\operatorname{End}_R(\mu_i(M))$ are derived equivalent, via the tilting $\operatorname{End}_R(M)$-module $\operatorname{Hom}_R(M, \mu_i(M))$.

We note that as in the above setup with R a Gorenstein normal 3-fold, a special case of 1.21 and 1.22 is, by 1.12, when $\operatorname{End}_R(M)$ is an arbitrary NCCR. Thus the above allows us to mutate any NCCR at any vertex, in particular the NCCR may be given by a quiver with relations where the quiver has both loops and 2-cycles. This situation happens very frequently in the study of one-dimensional fibres, where this form of mutation seems to have geometrical consequences.

We also point out that our mutation is a property of the algebra and not just the quiver. Under certain assumptions the quiver of the mutation is given geometrically in terms of some form of reconstruction, but note that in this level of generality a simple combinatorial rule (à la Fomin–Zelevinsky mutation) is too optimistic a hope. We do not mention quivers with potentials since we deal with algebras which in general may have infinite global dimension and so may have many more relations than arrows.

1.5. **Conventions.** We now state our conventions. All modules will be left modules, so for a ring A we denote $\operatorname{mod} A$ to be the category of finitely generated left A-modules. Throughout when composing maps fg will mean f then g, similarly for quivers ab will mean a then b. Note that with this convention $\operatorname{Hom}_R(M, X)$ is a $\operatorname{End}_R(M)$-module and $\operatorname{Hom}_R(X, M)$ is a $\operatorname{End}_R(M)^{\mathrm{op}}$-module. For $M \in \operatorname{mod} A$ we denote $\operatorname{add} M$ to be the full subcategory consisting of summands of finite direct sums of copies of M and we denote $\operatorname{proj} A := \operatorname{add} A$ to be the category of finitely generated projective A-modules. Throughout we will always use the letter R to denote some kind of *commutative noetherian* ring. We always strive to work in the global setting, so we write $(R, \mathfrak{m})$ if R is local.

2. Preliminaries

2.1. **Depth and CM Modules.** Here we record the preliminaries we shall need in subsequent sections, especially some global-local arguments that will be used extensively. For a commutative noetherian local ring $(R, \mathfrak{m})$ and $M \in \operatorname{mod} R$ recall that the *depth* of M is defined to be

$$\operatorname{depth}_R M := \inf\{i \geq 0 : \operatorname{Ext}^i_R(R/\mathfrak{m}, M) \neq 0\},$$

which coincides with the maximal length of a M-regular sequence. Keeping the assumption that $(R, \mathfrak{m})$ is local we say that $M \in \operatorname{mod} R$ is *maximal Cohen-Macaulay* (or simply, *CM*) if $\operatorname{depth}_R M = \dim R$. This definition generalizes to the non-local case as follows: if R is an arbitrary commutative noetherian ring we say that $M \in \operatorname{mod} R$ is *CM* if $M_{\mathfrak{p}}$ is CM for all prime ideals $\mathfrak{p}$ in R, and we say that R is a *CM ring* if R is a CM R-module.

It is often convenient to lift the CM property to noncommutative rings, which we do as follows:

Definition 2.1. *Let Λ be a module-finite R-algebra, then we call $M \in \operatorname{mod}\Lambda$ a* CM Λ-module *if it is CM when viewed as an R-module. We denote the category of CM Λ-modules by* $\operatorname{CM}\Lambda$.

To enable us to bring the concept of positive depth to non-local rings, the following is convenient:

Definition 2.2. *Let R be commutative, $M \in \operatorname{mod} R$. We denote $\mathsf{fl}\, M$ to be the unique maximal finite length R-submodule of M.*

It is clear that $\mathsf{fl}\, M$ exists because of the noetherian property of M; when $(R, \mathfrak{m})$ is local $\mathsf{fl}\, M = \{x \in M : \exists\, r \in \mathbb{N} \text{ with } \mathfrak{m}^r x = 0\}$. The following is well-known.

Lemma 2.3. *Let $(R, \mathfrak{m})$ be a local ring of dimension $d \geq 2$ and let Λ be a module finite R-algebra. Then for all $M, N \in \operatorname{mod}\Lambda$ with $\operatorname{depth}_R N \geq 2$ we have $\operatorname{depth}_R \operatorname{Hom}_\Lambda(M, N) \geq 2$.*

Proof. $\Lambda^a \to \Lambda^b \to M \to 0$ gives $0 \to \operatorname{Hom}_\Lambda(M, N) \to N^b \to N^a$ so the result follows from the depth lemma. □

In particular if $\operatorname{depth} R \geq 2$ then reflexive R-modules always have depth ≥ 2.

Lemma 2.4. *Suppose R is a d-dimensional CM ring with $d \geq 2$ and let Λ be a module finite R-algebra. For any $X \in \operatorname{ref}\Lambda$ we have $X_\mathfrak{p} \in \operatorname{CM}\Lambda_\mathfrak{p}$ for all $\mathfrak{p} \in \operatorname{Spec} R$ with $\operatorname{ht}\mathfrak{p} \leq 2$.*

Proof. Since X is reflexive as an R-module we can find an exact sequence $0 \to X \to P \to Q$ with $P, Q \in \operatorname{add} R$ and so on localizing we see that $X_\mathfrak{p}$ is a second syzygy for all primes $\mathfrak{p}$. Consequently if $\mathfrak{p}$ has height ≤ 2 then $X_\mathfrak{p}$ is a second syzygy for the CM ring $R_\mathfrak{p}$ which has $\dim R_\mathfrak{p} \leq 2$ and so $X_\mathfrak{p} \in \operatorname{CM} R_\mathfrak{p}$. □

2.2. **Reflexive Equivalence and Symmetric Algebras.** Here we introduce and fix notation for reflexive modules and symmetric algebras. All the material in this subsection can be found in [IR08]. Recall from the introduction (1.16) our convention on the definition of reflexive modules. Recall also that if Λ is a module finite R-algebra, we say $M \in \operatorname{ref}\Lambda$ is called a *height one progenerator* (respectively, *height one projective*) if $M_\mathfrak{p}$ is a progenerator (respectively, projective) over $\Lambda_\mathfrak{p}$ for all $\mathfrak{p} \in \operatorname{Spec} R$ with $\operatorname{ht}\mathfrak{p} \leq 1$.

In this paper, when the underlying commutative ring R is a normal domain the following reflexive equivalence is crucial:

Lemma 2.5. *If Λ is a module finite R-algebra, then*
(1) *If $M \in \operatorname{mod}\Lambda$ is a generator then $\operatorname{Hom}_\Lambda(M, -) : \operatorname{mod}\Lambda \to \operatorname{mod}\operatorname{End}_\Lambda(M)$ is fully faithful, restricting to an equivalence $\operatorname{add} M \xrightarrow{\approx} \operatorname{proj}\operatorname{End}_\Lambda(M)$.*
If further R is a normal domain, then the following assertions hold.
(2) *$\operatorname{Hom}_\Lambda(X, Y) \in \operatorname{ref} R$ for any $X \in \operatorname{mod}\Lambda$ and any $Y \in \operatorname{ref}\Lambda$.*
(3) *Every non-zero $M \in \operatorname{ref} R$ is a height one progenerator.*
(4) *Suppose Λ is a reflexive R-module and let $M \in \operatorname{ref}\Lambda$ be a height one progenerator. Then $\operatorname{Hom}_\Lambda(M, -) : \operatorname{ref}\Lambda \to \operatorname{ref}\operatorname{End}_\Lambda(M)$ is an equivalence. In particular $\operatorname{Hom}_R(N, -) : \operatorname{ref} R \to \operatorname{ref}\operatorname{End}_R(N)$ is an equivalence for all $N \in \operatorname{ref} R$.*

Proof. (1) is standard.
(2) is very easy; see [IR08, 2.4(1)].
(3) If $\mathfrak{p}$ is a height one prime then by 2.4 $M_\mathfrak{p} \in \operatorname{CM} R_\mathfrak{p}$. But R is normal so $R_\mathfrak{p}$ is regular; thus $M_\mathfrak{p}$ is free.
(4) follows by (3) and [RV89] (see also [IR08, 2.4(2)(i)]). □

The following lemma is convenient and will be used extensively.

Lemma 2.6. *Let R be a 3-dimensional, equi-codimensional CM ring and let Λ be a module finite R-algebra. If $X \in \operatorname{mod}\Lambda$ and $Y \in \operatorname{ref}\Lambda$ then*

$$\operatorname{Hom}_\Lambda(X, Y) \in \operatorname{CM} R \Rightarrow \mathsf{fl}\operatorname{Ext}^1_\Lambda(X, Y) = 0.$$

If further $Y \in \operatorname{CM}\Lambda$ then the converse holds.

Proof. ($\Rightarrow$) For each $\mathfrak{m} \in \operatorname{Max} R$ there is an exact sequence

$$0 \to \operatorname{Hom}_\Lambda(X,Y)_\mathfrak{m} \xrightarrow{f_\mathfrak{m}} \operatorname{Hom}_\Lambda(P,Y)_\mathfrak{m} \to \operatorname{Hom}_\Lambda(\Omega X,Y)_\mathfrak{m} \to \operatorname{Ext}^1_\Lambda(X,Y)_\mathfrak{m} \to 0 \quad (2.\mathrm{A})$$

obtained by localizing the exact sequence obtained from $0 \to \Omega X \to P \to X \to 0$ with $P \in \operatorname{add}\Lambda$. Now $\operatorname{depth}_{R_\mathfrak{m}} \operatorname{Hom}_{\Lambda_\mathfrak{m}}(X_\mathfrak{m}, Y_\mathfrak{m}) = 3$ and further by 2.3 $\operatorname{depth}_{R_\mathfrak{m}} \operatorname{Hom}_{\Lambda_\mathfrak{m}}(P_\mathfrak{m}, Y_\mathfrak{m}) \geq 2$, thus $\operatorname{depth}_{R_\mathfrak{m}} \operatorname{Cok} f_\mathfrak{m} \geq 2$. Since again by 2.3 $\operatorname{depth}_{R_\mathfrak{m}} \operatorname{Hom}_{\Lambda_\mathfrak{m}}(\Omega X_\mathfrak{m}, Y_\mathfrak{m}) \geq 2$, we conclude that $\operatorname{depth}_{R_\mathfrak{m}} \operatorname{Ext}^1_{\Lambda_\mathfrak{m}}(X_\mathfrak{m}, Y_\mathfrak{m}) > 0$ for all $\mathfrak{m} \in \operatorname{Max} R$ and so $\mathsf{fl}\operatorname{Ext}^1_\Lambda(X,Y) = 0$.
($\Leftarrow$) Suppose now that $Y \in \operatorname{CM}\Lambda$. Then in (2.A) $\operatorname{depth}_{R_\mathfrak{m}} \operatorname{Hom}_{\Lambda_\mathfrak{m}}(P_\mathfrak{m}, Y_\mathfrak{m}) = 3$ and so by a similar argument $\operatorname{depth}_{R_\mathfrak{m}} \operatorname{Ext}^1_{\Lambda_\mathfrak{m}}(X_\mathfrak{m}, Y_\mathfrak{m}) > 0$ implies that $\operatorname{depth}_{R_\mathfrak{m}} \operatorname{Hom}_{\Lambda_\mathfrak{m}}(X_\mathfrak{m}, Y_\mathfrak{m}) = 3$. ☐

Definition 2.7. *Let Λ be a module-finite R-algebra where R is an arbitrary commutative ring. We call Λ a* symmetric R-algebra *if $\operatorname{Hom}_R(\Lambda, R) \cong \Lambda$ as Λ-bimodules. We call Λ a* locally symmetric R-algebra *if $\Lambda_\mathfrak{p}$ is a symmetric $R_\mathfrak{p}$-algebra for all $\mathfrak{p} \in \operatorname{Spec} R$.*

Note that if Λ is a symmetric R-algebra then as functors $\operatorname{mod}\Lambda \to \operatorname{mod}\Lambda^{\mathrm{op}}$

$$\operatorname{Hom}_\Lambda(-,\Lambda) \cong \operatorname{Hom}_\Lambda(-, \operatorname{Hom}_R(\Lambda, R)) \cong \operatorname{Hom}_R(\Lambda \otimes_\Lambda -, R) = \operatorname{Hom}_R(-,R).$$

We have the following well-known observation.

Lemma 2.8. *Let R be a normal domain and Λ be a symmetric R-algebra. Then there is a functorial isomorphism $\operatorname{Hom}_\Lambda(X,Y) \cong \operatorname{Hom}_\Lambda(Y,X)^*$ for all $X, Y \in \operatorname{ref}\Lambda$ such that Y is height one projective.*

Proof. For the convenience of the reader, we give a detailed proof here. We have a natural map $f : \operatorname{Hom}_\Lambda(Y,\Lambda) \otimes_\Lambda X \to \operatorname{Hom}_\Lambda(Y,X)$ sending $a \otimes x$ to $(y \mapsto a(y)x)$. Consider the map $f^* : \operatorname{Hom}_\Lambda(Y,X)^* \to (\operatorname{Hom}_\Lambda(Y,\Lambda) \otimes_\Lambda X)^*$ between reflexive R-modules. Since Y is height one projective, $f_\mathfrak{p}$ and $(f^*)_\mathfrak{p}$ are isomorphisms for any prime $\mathfrak{p}$ of height at most one. Thus f^* is an isomorphism since R is normal. Thus we have

$$\operatorname{Hom}_\Lambda(Y,X)^* \overset{f^*}{\cong} (\operatorname{Hom}_\Lambda(Y,\Lambda) \otimes_\Lambda X)^* \cong (Y^* \otimes_\Lambda X)^* \cong \operatorname{Hom}_\Lambda(X, Y^{**}) \cong \operatorname{Hom}_\Lambda(X,Y)$$

as required. ☐

This immediately gives the following result, which implies that symmetric algebras are closed under reflexive equivalence.

Lemma 2.9. [IR08, 2.4(3)] *If Λ is a symmetric R-algebra then so is $\operatorname{End}_\Lambda(M)$ for any height one projective $M \in \operatorname{ref}\Lambda$. In particular if R is a normal domain and $N \in \operatorname{ref} R$ then $\operatorname{End}_R(N)$ is a symmetric R-algebra.*

2.3. Non-Singular and Gorenstein Orders.

Recall from 1.5 the definition of a non-singular R-order. By definition the localization of a non-singular R-order is again a non-singular R order — we shall see in 2.14 that in most situations we may check whether an algebra is a non-singular R-order by checking only at the maximal ideals.

For some examples of non-singular R-orders, recall that for a ring Λ and a finite group G together with a group homomorphism $G \to \operatorname{Aut}_{k-\mathrm{alg}}(\Lambda)$, we define the *skew group ring* $\Lambda \# G$ as follows [Aus86, Yos90]: As a set, it is a free Λ-module with the basis G. The multiplication is given by

$$(sg)(s'g') := (s \cdot g(s'))(gg')$$

for any $s, s' \in S$ and $g, g' \in G$.

This gives us a lot of examples of non-singular orders by the following result.

Lemma 2.10. *Let R be a CM ring containing a field k. Let Λ be a non-singular R-order, let G be a finite group together with a group homomorphism $G \to \operatorname{Aut}_{k-\mathrm{alg}}(\Lambda)$ and suppose $|G| \neq 0$ in k. Then $\Lambda \# G$ is a non-singular R-order.*

Proof. Since $\Lambda \# G$ is a direct sum of copies of Λ as an R-module and $\Lambda \in \operatorname{CM} R$, we have $\Lambda \# G \in \operatorname{CM} R$.

Let $X, Y \in \operatorname{mod}\Lambda \# G$. Then G acts on $\operatorname{Hom}_\Lambda(X,Y)$ by

$$(gf)(x) := g \cdot f(g^{-1}x)$$

for all $g \in G$, $f \in \mathrm{Hom}_\Lambda(X,Y)$ and $x \in X$. Clearly we have a functorial isomorphism

$$\mathrm{Hom}_{\Lambda\#G}(X,Y) = \mathrm{Hom}_\Lambda(X,Y)^G$$

for all $X, Y \in \mathrm{mod}\,\Lambda\#G$. The G-invariant $(-)^G$ is an exact functor since kG is semisimple. Thus we have a functorial isomorphism

$$\mathrm{Ext}^i_{\Lambda\#G}(X,Y) = \mathrm{Ext}^i_\Lambda(X,Y)^G$$

for all $X, Y \in \mathrm{mod}\,\Lambda\#G$ and $i \geq 0$. In particular, $\mathrm{gl.dim}\,\Lambda\#G \leq \mathrm{gl.dim}\,\Lambda$ holds, and we have the assertion. $\square$

Lemma 2.11. *Non-singular R-orders are closed under Morita equivalence.*

Proof. Suppose Λ is a non-singular R-order and Γ is Morita equivalent to Λ. Then $\Lambda_\mathfrak{p}$ is Morita equivalent to $\Gamma_\mathfrak{p}$ for all primes $\mathfrak{p}$. Thus since global dimension passes across Morita equivalence we have $\mathrm{gl.dim}\,\Gamma_\mathfrak{p} = \mathrm{gl.dim}\,\Lambda_\mathfrak{p} = \dim R_\mathfrak{p}$ for all primes $\mathfrak{p}$. To see why the CM property passes across the Morita equivalence let P denote the progenerator in $\mathrm{mod}\,\Lambda$ such that $\Gamma \cong \mathrm{End}_\Lambda(P)$. Since P is a summand of Λ^n for some n we know that Γ is a summand of $\mathrm{End}_\Lambda(\Lambda^n) = M_n(\Lambda)$ as an R-module. Since Λ is CM, so is Γ. $\square$

Recall from the introduction (§1.2) the definition of a canonical module ω_R for a non-local CM ring R. If Λ is an R-order we have an exact duality

$$\mathrm{Hom}_R(-,\omega_R) : \mathrm{CM}\,\Lambda \leftrightarrow \mathrm{CM}\,\Lambda^{\mathrm{op}}$$

and so the Λ-module $\omega_\Lambda := \mathrm{Hom}_R(\Lambda, \omega_R)$ is an injective cogenerator in the category $\mathrm{CM}\,\Lambda$.

Definition 2.12. [CR90, GN02] *Assume R has a canonical module ω_R. An R-order Λ is called* Gorenstein *if ω_Λ is a projective Λ-module.*

It is clear that if Λ is a Gorenstein R-order then $\Lambda_\mathfrak{p}$ is a Gorenstein $R_\mathfrak{p}$-order for all $\mathfrak{p} \in \mathrm{Spec}\,R$. Moreover one can show that Λ is Gorenstein $\iff$ $\mathrm{add}\,\Lambda = \mathrm{add}\,\omega_\Lambda \iff$ Λ^{op} is Gorenstein by using 4.11 and reducing to the complete local case. If Λ is 3-CY$^-$ (see §2.4) we shall see in 2.17 that Λ is a Gorenstein order.

When R is local, Gorenstein R-orders Λ are especially important since we have the following Auslander–Buchsbaum type equailty, which in particular says that the Λ-modules which are CM as R-modules are precisely the projective Λ-modules.

Lemma 2.13. *Let $(R, \mathfrak{m})$ be a local CM ring with a canonical module ω_R and let Λ be a Gorenstein R-order. Then for any $X \in \mathrm{mod}\,\Lambda$ with $\mathrm{proj.dim}_\Lambda X < \infty$,*

$$\mathrm{depth}_R\, X + \mathrm{proj.dim}_\Lambda\, X = \dim R.$$

Proof. Let X be a Λ-module with $\mathrm{proj.dim}_\Lambda X < \infty$.
(i) We will show that if $X \in \mathrm{CM}\,\Lambda$ then X is projective. We know that $\mathrm{Ext}^i_\Lambda(-, \omega_\Lambda) = 0$ on $\mathrm{CM}\,\Lambda$ for all $i > 0$ since ω_Λ is injective in $\mathrm{CM}\,\Lambda$. Since Λ is Gorenstein we have $\mathrm{Ext}^i_\Lambda(-, \Lambda) = 0$ on $\mathrm{CM}\,\Lambda$ for all $i > 0$. Since $\mathrm{Ext}^n_\Lambda(X, \Lambda) \neq 0$ for $n = \mathrm{proj.dim}\,X$, we have that X is projective.
(ii) Let $n = \mathrm{proj.dim}\,X$ and $t = \mathrm{depth}\,X$. Take a minimal projective resolution

$$0 \to P_n \to \ldots \to P_0 \to X \to 0.$$

By the depth lemma necessarily $t \geq d - n$. On the other hand by the depth lemma we have $\Omega^{d-t} X \in \mathrm{CM}\,\Lambda$. By (i) we know $\Omega^{d-t} X$ is projective so $n \leq d - t$. Thus $d = n + t$. $\square$

The following result is well-known to experts [Aus84].

Proposition 2.14. *Let Λ be an R-order where R is a CM ring with a canonical module ω_R. Then the following are equivalent:*
(1) *Λ is non-singular.*
(2) $\mathrm{gl.dim}\,\Lambda_\mathfrak{m} = \dim R_\mathfrak{m}$ *for all* $\mathfrak{m} \in \mathrm{Max}\,R$.
(3) $\mathrm{CM}\,\Lambda = \mathrm{proj}\,\Lambda$.
(4) *Λ is Gorenstein and* $\mathrm{gl.dim}\,\Lambda < \infty$.

Proof. (1)⇒(2) Immediate.
(2)⇒(3) Let $X \in \operatorname{CM}\Lambda$. Then $X_{\mathfrak{m}} \in \operatorname{CM}\Lambda_{\mathfrak{m}}$. Let $x_1, \dots, x_d$ be a $X_{\mathfrak{m}}$-regular sequence with $d = \dim R_{\mathfrak{m}}$. Then we have $\operatorname{proj.dim}_{\Lambda_{\mathfrak{m}}}(X_{\mathfrak{m}}/(x_1, \dots, x_d)X_{\mathfrak{m}}) = \operatorname{proj.dim}_{\Lambda_{\mathfrak{m}}} X_{\mathfrak{m}} + d$. This implies $X_{\mathfrak{m}}$ is a projective $\Lambda_{\mathfrak{m}}$-module for any $\mathfrak{m} \in \operatorname{Max} R$. Thus X is a projective Λ-module (see 4.11).
(3)⇒(4) We have $\omega_\Lambda \in \operatorname{CM}\Lambda = \operatorname{proj}\Lambda$. Since $\Omega^d X \in \operatorname{CM}\Lambda = \operatorname{proj}\Lambda$ for any $X \in \operatorname{mod}\Lambda$, we have $\operatorname{gl.dim}\Lambda \leq d$.
(4)⇒(1) Pick $\mathfrak{p} \in \operatorname{Spec} R$ and suppose $Y \in \operatorname{mod}\Lambda_{\mathfrak{p}}$. By Auslander–Buchsbaum 2.13 $\operatorname{proj.dim}_{\Lambda_{\mathfrak{p}}} Y \leq \dim R_{\mathfrak{p}}$ and so $\operatorname{gl.dim}\Lambda_{\mathfrak{p}} \leq \dim R_{\mathfrak{p}}$. Since $\Lambda_{\mathfrak{p}}$ is an $R_{\mathfrak{p}}$-order, $\operatorname{gl.dim}\Lambda_{\mathfrak{p}} \geq \dim R_{\mathfrak{p}}$. ☐

2.4. **d-CY$^-$ Algebras.** Throughout this paper we shall freely use the notion of d-CY and d-CY$^-$ as in [IR08, §3]: let R be a commutative noetherian ring with $\dim R = d$ and let Λ be a module-finite R-algebra. For any $X \in \operatorname{mod}\Lambda$ denote by $E(X)$ the injective hull of X, and put $E := \oplus_{\mathfrak{m}\in \operatorname{Max} R} E(R/\mathfrak{m})$. This gives rise to Matlis duality $D := \operatorname{Hom}_R(-, E)$ (see for example [O76, §1]). Matlis duality always gives a duality from the category of finite length R-modules to itself. This is true without assuming that R is (semi-)local because any finite length R-module is the finite direct sum of finite length $R_{\mathfrak{m}}$-modules for maximal ideals $\mathfrak{m}$, so the statement follows from that for the local setting [BH, 3.2.12].

Definition 2.15. *For $n \in \mathbb{Z}$ we call Λ n-CY if there is a functorial isomorphism*

$$\operatorname{Hom}_{\mathrm{D}(\operatorname{Mod}\Lambda)}(X, Y[n]) \cong D\operatorname{Hom}_{\mathrm{D}(\operatorname{Mod}\Lambda)}(Y, X)$$

for all $X, Y \in \mathrm{D}^{\mathrm{b}}(\mathsf{fl}\,\Lambda)$. Similarly we call Λ n-CY$^-$ if the above functorial isomorphism holds for any $X \in \mathrm{D}^{\mathrm{b}}(\mathsf{fl}\,\Lambda)$ and $Y \in \mathrm{K}^{\mathrm{b}}(\operatorname{proj}\Lambda)$.

We know [IR08, 3.1(7)] that Λ is n-CY if and only if it is n-CY$^-$ and $\operatorname{gl.dim}\Lambda < \infty$. The next two results can be found in [IR08]; we include both here since we use them extensively.

Proposition 2.16. [IR08, 3.10] *Let R be a commutative noetherian ring, $d \in \mathbb{N}$. Then R is d-CY$^-$ if and only if R is Gorenstein and equi-codimensional with $\dim R = d$.*

From this, for brevity we often say 'R is d-CY$^-$' instead of saying 'R is Gorenstein and equi-codimensional with $\dim R = d$'.

Proposition 2.17. *Let R be d-CY$^-$ and let Λ be a module-finite R-algebra. Then*

$$\Lambda \text{ is } d\text{-CY}^- \iff \Lambda \text{ is an } R\text{-order which is a locally symmetric } R\text{-algebra.}$$

Thus if Λ is d-CY$^-$ then Λ is a Gorenstein R-order.

Proof. The first statement is [IR08, 3.3(1)]. For the second, suppose Λ is d-CY$^-$ then since it is locally symmetric we have $\Lambda_{\mathfrak{m}} \cong \operatorname{Hom}_{R_{\mathfrak{m}}}(\Lambda_{\mathfrak{m}}, R_{\mathfrak{m}}) = \operatorname{Hom}_R(\Lambda, R)_{\mathfrak{m}}$ is a projective $\Lambda_{\mathfrak{m}}$-module for all $\mathfrak{m} \in \operatorname{Max} R$. Hence $\operatorname{Hom}_R(\Lambda, R)$ is a projective Λ-module, as required. ☐

The following picture for d-CY$^-$ rings R may help the reader navigate the terminology introduced above.

$$\begin{tikzcd}
d\text{-CY } R\text{-algebra} \arrow[rr, Rightarrow] \arrow[d, Rightarrow] & & \text{non-singular } R\text{-order} \arrow[d, Rightarrow, "2.14"] \\
d\text{-CY}^- \ R\text{-algebra} \arrow[r, Leftrightarrow, "2.17"] & \text{locally symmetric } R\text{-order} \arrow[r, Rightarrow, "2.17"] & \text{Gorenstein } R\text{-order}
\end{tikzcd}$$

The following non-local result is also useful.

Lemma 2.18. *Suppose R is a d-CY$^-$ normal domain.*
(1) If Λ is a module-finite R-algebra which is d-CY$^-$ and $M \in \operatorname{ref}\Lambda$ is a height one progenerator, then $\operatorname{End}_\Lambda(M)$ is d-CY$^-$ $\iff$ $\operatorname{End}_\Lambda(M) \in \operatorname{CM} R$.
(2) If $N \in \operatorname{ref} R$ then $\operatorname{End}_R(N)$ is d-CY$^-$ $\iff$ $\operatorname{End}_R(N) \in \operatorname{CM} R$.

Proof. (1) Let $\Gamma := \operatorname{End}_R(M)$. By 2.17 $\Lambda_{\mathfrak{m}}$ is a symmetric $R_{\mathfrak{m}}$-algebra for all $\mathfrak{m} \in \operatorname{Max} R$, thus $\Gamma_{\mathfrak{m}}$ is a symmetric $R_{\mathfrak{m}}$-algebra by 2.9. By 2.17, Γ is d-CY$^-$ if and only if $\Gamma_{\mathfrak{m}} \in \operatorname{CM} R_{\mathfrak{m}}$ for all $\mathfrak{m} \in \operatorname{Max} R$ if and only if $\Gamma \in \operatorname{CM} R$. Thus the assertion follows.
(2) This follows immediately from (1) since any $N \in \operatorname{ref} R$ is a height one progenerator by 2.5(3). ☐

Throughout we shall use the definition of NCCR in the introduction (1.6) due to its suitability for global-local arguments. We note that it is equivalent to the definition given by Van den Bergh:

Lemma 2.19. *Let R be a d-CY$^-$ normal domain, then $M \in \operatorname{ref} R$ gives a NCCR if and only if* $\operatorname{gl.dim} \operatorname{End}_R(M) < \infty$ *and* $\operatorname{End}_R(M) \in \operatorname{CM} R$.

Proof. ($\Rightarrow$) obvious.
($\Leftarrow$) Set $\Lambda := \operatorname{End}_R(M)$, $d := \dim R$. By 2.18(2) Λ is d-CY$^-$ hence by 2.17 Λ is a Gorenstein order, with $\operatorname{gl.dim} \Lambda < \infty$. By 2.14 Λ is non-singular. $\square$

3. AUSLANDER–REITEN DUALITY FOR NON-ISOLATED SINGULARITIES

Let R be a d-dimensional, equi-codimensional CM ring with a canonical module ω_R, and let Λ be an R-order. We denote $\underline{\mathrm{CM}}\Lambda$ to be the stable category of CM Λ-modules and $\overline{\mathrm{CM}}\Lambda$ to be the costable category. We have *AR translation*

$$\tau := \operatorname{Hom}_R(\Omega^d_{\Lambda^{\mathrm{op}}} \operatorname{Tr}_\Lambda(-), \omega_R) : \underline{\mathrm{CM}}\Lambda \to \overline{\mathrm{CM}}\Lambda.$$

If further R is an isolated singularity then by Auslander–Reiten duality there is a functorial isomorphism

$$\underline{\operatorname{Hom}}_R(X, Y) \cong D \operatorname{Ext}^1_R(Y, \tau X)$$

for all $X, Y \in \operatorname{CM} R$, where D is the Matlis dual (as in §2.4). However, in what follows we do not assume that R is an isolated singularity.

If Λ is an R-order as above we define $\operatorname{Sing} \Lambda := \{\mathfrak{p} \in \operatorname{Spec} R : \Lambda_{\mathfrak{p}} \text{ is singular}\}$ to be the singular locus of Λ. Our main theorem is the following:

Theorem 3.1. *Let R be a d-dimensional, equi-codimensional CM ring with a canonical module ω_R. Let Λ be an R-order with* $\dim \operatorname{Sing} \Lambda \leq 1$. *Then there exists a functorial isomorphism*

$$\mathsf{fl}\, \underline{\operatorname{Hom}}_\Lambda(X, Y) \cong D(\mathsf{fl} \operatorname{Ext}^1_\Lambda(Y, \tau X))$$

for all $X, Y \in \operatorname{CM} \Lambda$.

For normal 3-folds, the key consequence is 3.8(2). In fact 3.1 immediately follows from the more general 3.2 below. Recall for $X \in \operatorname{mod} \Lambda$ that $\operatorname{NP}(X) := \{\mathfrak{p} \in \operatorname{Spec} R : X_{\mathfrak{p}} \text{ is non-projective}\}$ and $\operatorname{CM}_1 \Lambda := \{X \in \operatorname{CM} \Lambda : \dim \operatorname{NP}(X) \leq 1\}$.

Theorem 3.2. *Let R be a d-dimensional, equi-codimensional CM ring with a canonical module ω_R. Let Λ be an R-order. Then there exists a functorial isomorphism*

$$\mathsf{fl}\, \underline{\operatorname{Hom}}_\Lambda(X, Y) \cong D(\mathsf{fl} \operatorname{Ext}^1_\Lambda(Y, \tau X))$$

for all $X \in \operatorname{CM}_1 \Lambda$ *and* $Y \in \operatorname{CM} \Lambda$

The proof of 3.2 requires the next four easy lemmas. For a finitely generated R-module M, denote $E_R(M)$ to be the injective hull of M.

Lemma 3.3. *If $X \in \operatorname{mod} R$ and $Y \in \operatorname{Mod} R$ satisfies* $\operatorname{Supp} X \cap \operatorname{Ass} Y = \emptyset$, *then* $\operatorname{Hom}_R(X, Y) = 0$.

Proof. Let $f : X \to Y$ be any map and $X' := \operatorname{Im} f$. Then $X' \subset Y$ is a finitely generated submodule such that $\operatorname{Ass} X' \subset \operatorname{Supp} X \cap \operatorname{Ass} Y$. Thus $\operatorname{Ass} X' = \emptyset$ and so since X' is finitely generated, $X' = 0$. $\square$

Lemma 3.4. (1) *We have $E_R(R/\mathfrak{p}) = E_R(R/\mathfrak{p})_{\mathfrak{p}}$ as R-modules, and this is isomorphic to $E_{R_{\mathfrak{p}}}(R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}})$ as $R_{\mathfrak{p}}$-modules (so also as R-modules).*
(2) $\operatorname{Ass} E_R(R/\mathfrak{p}) = \{\mathfrak{p}\}$.

Proof. (1) The former assertion is [Lam, 3.77], and the latter assertion is [BH, 3.2.5].
(2) This is [BH, 3.2.7(a)]. $\square$

In what follows we often write $E(M)$ instead of $E_R(M)$.

Lemma 3.5. *If* $\dim R/\mathfrak{p} \leq \dim R/\mathfrak{q}$ *then* $\operatorname{Hom}_R(R/\mathfrak{p}, E(R/\mathfrak{q})) = \left\{ \begin{array}{cc} 0 & \mathfrak{p} \neq \mathfrak{q} \\ R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}} & \mathfrak{p} = \mathfrak{q} \end{array} \right.$.

Proof. Assume $\mathfrak{p} \neq \mathfrak{q}$. Then $\mathfrak{q} \notin \operatorname{Supp} R/\mathfrak{p}$ and we have $\operatorname{Supp} R/\mathfrak{p} \cap \operatorname{Ass} E(R/\mathfrak{q}) = \emptyset$ by 3.4(2). Thus we have $\operatorname{Hom}_R(R/\mathfrak{p}, E(R/\mathfrak{q})) = 0$ by 3.3.

If $\mathfrak{p} = \mathfrak{q}$ then by change of rings

$$\operatorname{Hom}_R(R/\mathfrak{p}, E(R/\mathfrak{p})) \overset{3.4}{=} \operatorname{Hom}_R(R/\mathfrak{p}, E(R/\mathfrak{p})_{\mathfrak{p}}) = \operatorname{Hom}_{R_{\mathfrak{p}}}(R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}}, E(R/\mathfrak{p})_{\mathfrak{p}}) = R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}}$$

where the last equality is just [BH, 3.2.7(b)]. □

Lemma 3.6. *Let R be a commutative noetherian ring and X be a finitely generated R-module. Then we have a bijection between finite length submodules of X and finite length factor modules of DX, which is given by $Y \mapsto DY$.*

Proof. Matlis duality $D = \operatorname{Hom}_R(-, E)$ with $E = \oplus_{\mathfrak{m} \in \operatorname{Max} R} E(R/\mathfrak{m})$ gives a duality from the category of finite length modules to itself. Thus the assertion is valid if X has finite length, and in general the map is well-defined and injective. We only have to show the surjectivity.

Let Z be a finite length factor module of DX. For the ideal $I := \operatorname{Ann}_R Z$, we have that R/I is a finite length R-module. Now $X' := \operatorname{Hom}_R(R/I, X)$ is a finite length submodule of X, and we have

$$DX' = \operatorname{Hom}_R(\operatorname{Hom}_R(R/I, X), E)) \cong R/I \otimes_R \operatorname{Hom}_R(X, E) = R/I \otimes_R DX.$$

Thus Z is a factor module of DX' from our choice of I. Since X' has finite length, there exists a finite length submodule Y of X' (hence of X) such that $DY = Z$. □

We are now ready to prove 3.2:

Proof. Denote $T := \operatorname{Tr} X$. Consider the minimal R-injective resolution

$$0 \to \omega_R \to I_0 \to I_1 \to \ldots \to I_{d-1} \to I_d \to 0$$

of ω_R, then

$$I_i \overset{[\text{BH, 3.2.9, 3.3.10(b)}]}{=} \bigoplus_{\mathfrak{p}:\operatorname{ht}\mathfrak{p}=i} E(R/\mathfrak{p}) \overset{1.3}{=} \bigoplus_{\mathfrak{p}:\dim R/\mathfrak{p}=d-i} E(R/\mathfrak{p}). \tag{3.B}$$

In particular the Matlis dual $D = \operatorname{Hom}_R(-, I_d)$. Now since $Y \in \operatorname{CM}\Lambda$ we have $\operatorname{Ext}^i_R(Y, \omega_R) = 0$ for all $i > 0$ and so applying $\operatorname{Hom}_R(Y, -)$ gives an exact sequence

$$0 \to {}_R(Y, \omega_R) \to {}_R(Y, I_0) \to {}_R(Y, I_1) \to \ldots \to {}_R(Y, I_{d-1}) \to {}_R(Y, I_d) \to 0$$

of left Λ^{op}-modules, which we split into short exact sequences as

$$\begin{tikzcd}
0 \arrow[r] & {}_R(Y,\omega_R) \arrow[r] & {}_R(Y,I_0) \arrow[r] \arrow[d, two heads] & {}_R(Y,I_1) \arrow[r] \arrow[d, two heads] & {}_R(Y,I_2) \arrow[r] & \ldots \arrow[r] & {}_R(Y,I_{d-2}) \arrow[r] \arrow[d, two heads] & {}_R(Y,I_{d-1}) \arrow[r] & {}_R(Y,I_d) \arrow[r] & 0\,. \\
 & & C_1 \arrow[ur, hook] & C_2 \arrow[ur, hook] & & & C_{d-1} \arrow[ur, hook] & & &
\end{tikzcd}$$

Applying $\operatorname{Hom}_{\Lambda^{\mathrm{op}}}(T, -)$ gives exact sequences

$$\begin{aligned}
&\operatorname{Ext}^1_{\Lambda^{\mathrm{op}}}(T, {}_R(Y, I_{d-1})) \to \operatorname{Ext}^1_{\Lambda^{\mathrm{op}}}(T, {}_R(Y, I_d)) \longrightarrow \operatorname{Ext}^2_{\Lambda^{\mathrm{op}}}(T, C_{d-1}) \longrightarrow \operatorname{Ext}^2_{\Lambda^{\mathrm{op}}}(T, {}_R(Y, I_{d-1})) \\
&\operatorname{Ext}^2_{\Lambda^{\mathrm{op}}}(T, {}_R(Y, I_{d-2})) \longrightarrow \operatorname{Ext}^2_{\Lambda^{\mathrm{op}}}(T, C_{d-1}) \longrightarrow \operatorname{Ext}^3_{\Lambda^{\mathrm{op}}}(T, C_{d-2}) \longrightarrow \operatorname{Ext}^3_{\Lambda^{\mathrm{op}}}(T, {}_R(Y, I_{d-2})) \\
&\qquad\qquad\qquad\qquad\vdots \\
&\operatorname{Ext}^{d-1}_{\Lambda^{\mathrm{op}}}(T, {}_R(Y, I_1)) \longrightarrow \operatorname{Ext}^{d-1}_{\Lambda^{\mathrm{op}}}(T, C_2) \longrightarrow \operatorname{Ext}^d_{\Lambda^{\mathrm{op}}}(T, C_1) \longrightarrow \operatorname{Ext}^d_{\Lambda^{\mathrm{op}}}(T, {}_R(Y, I_1)) \\
&\operatorname{Ext}^d_{\Lambda^{\mathrm{op}}}(T, {}_R(Y, I_0)) \longrightarrow \operatorname{Ext}^d_{\Lambda^{\mathrm{op}}}(T, C_1) \longrightarrow \operatorname{Ext}^{d+1}_{\Lambda^{\mathrm{op}}}(T, {}_R(Y, \omega_R)) \longrightarrow \operatorname{Ext}^{d+1}_{\Lambda^{\mathrm{op}}}(T, {}_R(Y, I_0))\,.
\end{aligned}$$

By the assumption that $X \in \operatorname{CM}_1 \Lambda$, for all primes $\mathfrak{p}$ such that $\dim R/\mathfrak{p} > 1$, we have $X_{\mathfrak{p}} \in \operatorname{proj} \Lambda_{\mathfrak{p}}$ and so $T_{\mathfrak{p}} \in \operatorname{proj} \Lambda^{\mathrm{op}}_{\mathfrak{p}}$. Thus for all such primes and any $j > 0$, we have $\operatorname{Tor}^{\Lambda}_j(T, Y)_{\mathfrak{p}} \cong \operatorname{Tor}^{\Lambda_{\mathfrak{p}}}_j(T_{\mathfrak{p}}, Y_{\mathfrak{p}}) = 0$. Hence for all $i = 0, 1, \ldots, d-2$ and all $j > 0$, by 3.4(2) and (3.B) it follows that $\operatorname{Supp} \operatorname{Tor}^{\Lambda}_j(T, Y) \cap \operatorname{Ass} I_i = \emptyset$ and so consequently $\operatorname{Hom}_R(\operatorname{Tor}^{\Lambda}_j(T, Y), I_i) = 0$ for all $j > 0$ and all $i = 0, 1, \ldots, d-2$ by 3.3.

But using this vanishing of Hom together with the functorial isomorphism

$$\operatorname{Ext}^j_{\Lambda}(A, {}_R(B, I)) \cong \operatorname{Hom}_R(\operatorname{Tor}^{\Lambda}_j(A, B), I)$$

[CE99] (where I is an injective R-module) the above exact sequences reduce to

$${}_R(\mathrm{Tor}_1^\Lambda(T,Y),I_{d-1}) \longrightarrow {}_R(\mathrm{Tor}_1^\Lambda(T,Y),I_d) \longrightarrow \mathrm{Ext}^2_{\Lambda^{\mathrm{op}}}(T,C_{d-1}) \longrightarrow {}_R(\mathrm{Tor}_2^\Lambda(T,Y),I_{d-1})$$

with $\mathrm{Ext}^2_{\Lambda^{\mathrm{op}}}(T,C_{d-1}) \cong \mathrm{Ext}^3_{\Lambda^{\mathrm{op}}}(T,C_{d-2}) \cong \ldots \cong \mathrm{Ext}^d_{\Lambda^{\mathrm{op}}}(T,C_1) \cong \mathrm{Ext}^{d+1}_{\Lambda^{\mathrm{op}}}(T,{}_R(Y,\omega_R))$. Thus there is an exact sequence

$$\begin{tikzcd}
{}_R(\mathrm{Tor}_1^\Lambda(T,Y),I_{d-1}) \arrow[r] & {}_R(\mathrm{Tor}_1^\Lambda(T,Y),I_d) \arrow[r] \arrow[d, "\cong"'] & \mathrm{Ext}^{d+1}_{\Lambda^{\mathrm{op}}}(T,{}_R(Y,\omega_R)) \arrow[r] \arrow[d, "\cong"'] & {}_R(\mathrm{Tor}_2^\Lambda(T,Y),I_{d-1}) \\
 & D(\underline{\mathrm{Hom}}_\Lambda(X,Y)) \arrow[r, "\psi"] & \mathrm{Ext}^1_\Lambda(Y,\tau X) &
\end{tikzcd}$$

where the left hand isomorphism follows from the well-known fact that $\mathrm{Tor}_1^\Lambda(\mathrm{Tr}\,X,Y) \cong \underline{\mathrm{Hom}}_\Lambda(X,Y)$ [Yos90], and the right hand isomorphism from

$$\begin{aligned}\mathrm{Ext}^{d+1}_{\Lambda^{\mathrm{op}}}(T,{}_R(Y,\omega_R)) \cong \mathrm{Ext}^1_{\Lambda^{\mathrm{op}}}(\Omega^d_{\Lambda^{\mathrm{op}}}\mathrm{Tr}\,X,{}_R(Y,\omega_R)) &\cong \mathrm{Ext}^1_\Lambda(Y,{}_R(\Omega^d_{\Lambda^{\mathrm{op}}}\mathrm{Tr}\,X,\omega_R)) \\ &= \mathrm{Ext}^1_\Lambda(Y,\tau X).\end{aligned}$$

We claim that $\mathrm{Im}\,\psi$ has finite length. We know that $\mathrm{Im}\,\psi$ is noetherian. If $\mathrm{Im}\,\psi$ has infinite length then there exists an infinite sequence $\ldots \twoheadrightarrow Y_3 \twoheadrightarrow Y_2 \twoheadrightarrow Y_1$ of finite length factor modules of $\mathrm{Im}\,\psi$. By 3.6, we have an infinite ascending chain of submodules of $\underline{\mathrm{Hom}}_\Lambda(X,Y)$, a contradiction. Hence $\mathrm{Im}\,\psi$ has finite length.

We now claim that $\mathsf{fl}\,\mathrm{Hom}_R(\mathrm{Tor}_2^\Lambda(T,Y),I_{d-1})=0$, from which $\mathrm{Im}\,\psi = \mathsf{fl}\,\mathrm{Ext}^1_\Lambda(Y,\tau X)$ follows. But this is easy: take a surjection $R^n \twoheadrightarrow \mathrm{Tor}_2^\Lambda(T,Y)$ of R-modules and apply $\mathrm{Hom}_R(-,I_{d-1})$ to obtain $0 \to \mathrm{Hom}_R(\mathrm{Tor}_2^\Lambda(T,Y),I_{d-1}) \to I_{d-1}^n$. Since $\mathfrak{m} \notin \mathrm{Ass}\,I_{d-1}$ by 3.4(2) we have that I_{d-1} does not contain any finite length submodules and so the same is true of $\mathrm{Hom}_R(\mathrm{Tor}_2^\Lambda(T,Y),I_{d-1})$.

Finally we claim that $\mathrm{Hom}_R(\mathrm{Tor}_1^\Lambda(T,Y),I_{d-1})$ has no non-zero factor modules with finite length, as this then implies that $\mathrm{Ker}\,\psi$ has no non-zero factor modules with finite length and so $\mathrm{Im}\,\psi = D(\mathsf{fl}\,\underline{\mathrm{Hom}}_\Lambda(X,Y))$, completing the proof.

By the above $\dim_R \mathrm{Tor}_1^\Lambda(T,Y) \leq 1$ so $\mathrm{Tor}_1^\Lambda(T,Y)$ is filtered by some $R/\mathfrak{p}$ with $\dim R/\mathfrak{p} \leq 1$. It follows that $\mathrm{Hom}_R(\mathrm{Tor}_1^\Lambda(T,Y),I_{d-1})$ is filtered by some $\mathrm{Hom}_R(R/\mathfrak{p},I_{d-1})$ with $\dim R/\mathfrak{p} \leq 1$. But by 3.5 $\mathrm{Hom}_R(R/\mathfrak{p},I_{d-1}) \cong R_\mathfrak{p}/\mathfrak{p}R_\mathfrak{p}$ when $\dim R/\mathfrak{p}=1$ and $\mathrm{Hom}_R(R/\mathfrak{p},I_{d-1}) \cong 0$ when $\dim R/\mathfrak{p}=0$, which both have no non-zero factor modules of finite length. Thus $\mathrm{Hom}_R(\mathrm{Tor}_1^\Lambda(T,Y),I_{d-1})$ has no non-zero factor modules of finite length. $\square$

When R has only isolated singularities the above reduces to classical Auslander–Reiten duality. If moreover R is a 3-CY$^-$ ring with isolated singularities (i.e. R is a Gorenstein 3-dimensional equi-codimensional ring with isolated singularities), AR duality implies that the category $\underline{\mathrm{CM}}R$ is 2-CY. We now apply 3.1 to possibly non-isolated 3-CY$^-$ rings to obtain some analogue of this 2-CY property (see 3.8(1) below). The following lemma is well-known:

Lemma 3.7. *Suppose R is d-CY$^-$ and let Λ be a symmetric R-order. Then $\tau \cong \Omega_\Lambda^{2-d}$, thus for 3-folds $\tau^{-1} \cong \Omega_\Lambda$.*

Proof. We have $\Omega^2\,\mathrm{Tr}(-) \cong \mathrm{Hom}_\Lambda(-,\Lambda)$. Since R is Gorenstein and Λ is symmetric, we have $\Omega^2\,\mathrm{Tr}(-) \cong \mathrm{Hom}_\Lambda(-,\Lambda) \cong \mathrm{Hom}_R(-,\omega_R)$. Thus

$$\begin{aligned}\tau = \mathrm{Hom}_R(\Omega^d_{\Lambda^{\mathrm{op}}}\mathrm{Tr}(-),\omega_R) &\cong \mathrm{Hom}_R(\Omega^{d-2}_{\Lambda^{\mathrm{op}}}\mathrm{Hom}_R(-,\omega_R),\omega_R) \\ &\cong \Omega_\Lambda^{2-d}\,\mathrm{Hom}_R(\mathrm{Hom}_R(-,\omega_R),\omega_R) \\ &\cong \Omega_\Lambda^{2-d}.\end{aligned}$$

$\square$

Corollary 3.8. *Let R be 3-CY$^-$ and let Λ be a symmetric R-order with $\dim \mathrm{Sing}\,\Lambda \leq 1$. Then*

(1) *There is a functorial isomorphism*

$$\mathsf{fl}\,\mathrm{Ext}^1_\Lambda(X,Y) \cong D(\mathsf{fl}\,\mathrm{Ext}^1_\Lambda(Y,X))$$

for all $X, Y \in \mathrm{CM}\,\Lambda$.
(2) *If* $X, Y \in \mathrm{CM}\,\Lambda$ *then*

$$\mathrm{Hom}_\Lambda(X,Y) \in \mathrm{CM}\,R \Leftrightarrow \mathsf{fl}\,\mathrm{Ext}^1_\Lambda(X,Y) = 0 \Leftrightarrow \mathsf{fl}\,\mathrm{Ext}^1_\Lambda(Y,X) = 0 \Leftrightarrow \mathrm{Hom}_\Lambda(Y,X) \in \mathrm{CM}\,R.$$

Proof. (1) $D(\mathsf{fl}\,\mathrm{Ext}^1_\Lambda(Y,X)) \overset{3.1}{\cong} \mathsf{fl}\,\underline{\mathrm{Hom}}_\Lambda(\tau^{-1}X,Y) \overset{3.7}{\cong} \mathsf{fl}\,\underline{\mathrm{Hom}}_\Lambda(\Omega_\Lambda X,Y) \cong \mathsf{fl}\,\mathrm{Ext}^1_\Lambda(X,Y)$.
(2) Immediate from (1) and 2.6. □

Note that 3.8(2) also holds in greater generality by 2.8, but does not hold in the more general setting of CM rings.

4. Modifying and Maximal Modifying Modules

Motivated by the fact that $\mathrm{Spec}\,R$ need not have a crepant resolution, we want to be able to control infinite global dimension algebras and hence partial resolutions of singularities. We begin with our main definition.

Definition 4.1. *Let R be a d-dimensional CM ring, Λ a module finite R-algebra. We call $N \in \mathrm{ref}\,\Lambda$ a* modifying module *if $\mathrm{End}_\Lambda(N) \in \mathrm{CM}\,R$, whereas we call N a* maximal modifying (MM) module *if N is modifying and furthermore it is maximal with respect to this property, that is to say if there exists $X \in \mathrm{ref}\,\Lambda$ with $N \oplus X$ modifying, necessarily $X \in \mathrm{add}\,N$.*

The following is trivial but worth recording:

Lemma 4.2. *Suppose R is a d-dimensional CM ring, Λ a module finite R-algebra. Then*
(1) The modifying Λ-modules which are generators are always CM.
(2) If further R and Λ are both d-CY$^-$ then the MM generators are precisely the MM modules which are CM.

Proof. (1) If M is a modifying generator then $\mathrm{End}_\Lambda(M) \in \mathrm{CM}\,R$ so $M \cong \mathrm{Hom}_\Lambda(\Lambda, M) \in \mathrm{CM}\,R$ since it is a summand of $\mathrm{End}_\Lambda(M)$.
(2) Conversely suppose that M is a MM module which is CM. We have $\mathrm{End}_\Lambda(\Lambda) \in \mathrm{CM}\,R$ by 2.17. Further R is now Gorenstein (by 2.16) and Λ is locally symmetric (by 2.17), thus since $M \in \mathrm{CM}\,R$ we have

$$\mathrm{Hom}_{\Lambda_\mathfrak{m}}(M_\mathfrak{m}, \Lambda_\mathfrak{m}) \cong \mathrm{Hom}_{R_\mathfrak{m}}(M_\mathfrak{m}, R_\mathfrak{m}) \in \mathrm{CM}\,R_\mathfrak{m}$$

for all $\mathfrak{m} \in \mathrm{Max}\,R$ and so $\mathrm{Hom}_\Lambda(M,\Lambda) \in \mathrm{CM}\,R$. Trivially $M \cong \mathrm{Hom}_\Lambda(\Lambda, M) \in \mathrm{CM}\,R$. Hence $\mathrm{End}_\Lambda(M \oplus \Lambda) \in \mathrm{CM}\,R$ and so since M is maximal necessarily $\Lambda \in \mathrm{add}\,M$. □

Leading up to our next proposition (4.6) we require three useful technical lemmas:

Lemma 4.3. *Let R be a normal 3-CY$^-$ ring and let Λ be a module finite R-algebra which is 3-CY$^-$. Let $B \in \mathrm{ref}\,\Lambda$ be a modifying height one progenerator and let $C \in \mathrm{ref}\,\Lambda$. If $0 \to A \xrightarrow{f} B_0 \xrightarrow{g} C \to 0$ is an exact sequence where g is a right $\mathrm{add}\,B$-approximation, then the cokernel of the natural map*

$$\mathrm{Hom}_\Lambda(B_0, B) \xrightarrow{f\cdot} \mathrm{Hom}_\Lambda(A,B)$$

has finite length.

Proof. Set $\Gamma := \mathrm{End}_\Lambda(B)$. Since B is a height one progenerator we have a reflexive equivalence $\mathbb{F} := \mathrm{Hom}_\Lambda(B,-) : \mathrm{ref}\,\Lambda \to \mathrm{ref}\,\Gamma$ by 2.5(4). Moreover Γ is 3-CY$^-$ by 2.18.

Consider the exact sequence

$$0 \to \mathbb{F}A \to \mathbb{F}B_0 \to \mathbb{F}C \to 0$$

of Γ-modules. Then since $\mathbb{F}B_0 \in \mathrm{proj}\,\Gamma$, applying $\mathrm{Hom}_\Gamma(-,\Gamma) = \mathrm{Hom}_\Gamma(-,\mathbb{F}B)$ gives an exact sequence

$$\begin{tikzcd}
\mathrm{Hom}_\Gamma(\mathbb{F}B_0,\mathbb{F}B) \arrow[r] & \mathrm{Hom}_\Gamma(\mathbb{F}A,\mathbb{F}B) \arrow[r] & \mathrm{Ext}^1_\Gamma(\mathbb{F}C,\Gamma) \arrow[r] & 0 \\
\mathrm{Hom}_\Lambda(B_0,B) \arrow[u, "\cong"] \arrow[r, "f\cdot"] & \mathrm{Hom}_\Lambda(A,B) \arrow[u, "\cong"] & &
\end{tikzcd}$$

and thus $\mathrm{Cok}(f\cdot) = \mathrm{Ext}^1_\Gamma(\mathbb{F}C,\Gamma)$. Hence we only have to show that $\mathrm{Ext}^1_\Gamma(\mathbb{F}C,\Gamma)_\mathfrak{p} = 0$ for any non-maximal prime ideal $\mathfrak{p}$ of R. By 2.3 and 2.4 we have $(\mathbb{F}C)_\mathfrak{p} \in \mathrm{CM}\,\Gamma_\mathfrak{p}$.

Since Γ is 3-CY$^-$, $\Gamma_{\mathfrak{p}}$ is a Gorenstein $R_{\mathfrak{p}}$-order by 2.17. Consequently $\operatorname{Ext}^1_\Gamma(\mathbb{F}C,\Gamma)_{\mathfrak{p}} = \operatorname{Ext}^1_{\Gamma_{\mathfrak{p}}}((\mathbb{F}C)_{\mathfrak{p}},\Gamma_{\mathfrak{p}}) = 0$, as required. ☐

Lemma 4.4. *Let R be a normal 3-CY$^-$ ring and let Λ be a module finite R-algebra which is 3-CY$^-$. Suppose $N \in \operatorname{ref}\Lambda$ and $M \in \operatorname{CM}\Lambda$ with both M and N modifying such that M is a height one progenerator. If $0 \to L \to M_0 \xrightarrow{h} N \to 0$ is an exact sequence where h is a right $\operatorname{add} M$-approximation, then $\operatorname{End}_\Lambda(L \oplus M) \in \operatorname{CM} R$.*

Proof. Note first that since N is reflexive and $M \in \operatorname{CM}\Lambda$ we have $L \in \operatorname{CM}\Lambda$ by the depth lemma. From the exact sequence

$$0 \to \operatorname{Hom}_\Lambda(M,L) \to \operatorname{Hom}_\Lambda(M,M_0) \to \operatorname{Hom}_\Lambda(M,N) \to 0$$

with $\operatorname{Hom}_\Lambda(M,M_0) \in \operatorname{CM} R$ we see, using 2.3 and the depth lemma, that $\operatorname{Hom}_\Lambda(M,L) \in \operatorname{CM} R$. By 2.8 $\operatorname{Hom}_\Lambda(L,M) \in \operatorname{CM} R$. Since $\operatorname{End}_\Lambda(M) \in \operatorname{CM} R$ by assumption, it suffices to show that $\operatorname{End}_\Lambda(L) \in \operatorname{CM} R$. By 2.6 we only need to show that $\operatorname{fl}\operatorname{Ext}^1_\Lambda(L,L) = 0$.

Consider now the following exact commutative diagram

$$\begin{tikzcd}
 & & \operatorname{Cok} f \arrow[dr, hook] & & & \\
\operatorname{Hom}_\Lambda(L,M_0) \arrow[r, "f"] & \operatorname{Hom}_\Lambda(L,N) \arrow[ur, two heads] \arrow[rr] & & \operatorname{Ext}^1_\Lambda(L,L) \arrow[rr] \arrow[dr, two heads] & & \operatorname{Ext}^1_\Lambda(L,M_0) . \\
\operatorname{Hom}_\Lambda(M_0,M_0) \arrow[u, "b"] \arrow[r, two heads, "t"] & \operatorname{Hom}_\Lambda(M_0,N) \arrow[u, "c"] & & & K \arrow[ur, hook] &
\end{tikzcd}$$

Since $\operatorname{Hom}_\Lambda(L,M) \in \operatorname{CM} R$ we know that $\operatorname{fl}\operatorname{Ext}^1_\Lambda(L,M_0) = 0$ and so $\operatorname{fl} K = 0$. Hence to show that $\operatorname{fl}\operatorname{Ext}^1_\Lambda(L,L) = 0$ we just need to show that $\operatorname{fl}\operatorname{Cok} f = 0$. To do this consider the exact sequence

$$\operatorname{Cok} b \to \operatorname{Cok} bf \to \operatorname{Cok} f \to 0. \tag{4.C}$$

By 4.3 applied with $B = M_0$, $\operatorname{Cok} b$ has finite length and thus the image of the first map in (4.C) has finite length. Secondly note that $\operatorname{Cok} bf = \operatorname{Cok} c$ and $\operatorname{fl}\operatorname{Cok} c = 0$ since $\operatorname{Cok} c$ embeds inside $\operatorname{Ext}^1_\Lambda(N,N)$ and furthermore $\operatorname{fl}\operatorname{Ext}^1_\Lambda(N,N) = 0$ by 2.6. This means that the image of the first map is zero, hence $\operatorname{Cok} f \cong \operatorname{Cok} c$ and so in particular $\operatorname{fl}\operatorname{Cok} f = 0$. ☐

In fact, using reflexive equivalence we have the following improvement of 4.4 which does not assume that M is CM:

Lemma 4.5. *Let R be normal 3-CY$^-$ and let M and N be modifying modules. If $0 \to L \to M_0 \xrightarrow{h} N$ is an exact sequence where h is a right $\operatorname{add} M$-approximation, then $L \oplus M$ is modifying.*

Proof. Note that L is reflexive since R is normal. Denote $\Lambda := \operatorname{End}_R(M)$ and $\mathbb{F} := \operatorname{Hom}_R(M,-) : \operatorname{ref} R \to \operatorname{ref}\Lambda$ the reflexive equivalence in 2.5(4). Then Λ is 3-CY$^-$ by 2.18, $\mathbb{F}N \in \operatorname{ref}\Lambda$, $\mathbb{F}M \in \operatorname{CM}\Lambda$ and both $\mathbb{F}N$ and $\mathbb{F}M$ are modifying Λ-modules. Further

$$0 \to \mathbb{F}L \to \mathbb{F}M_0 \to \mathbb{F}N \to 0$$

is exact and trivially $\mathbb{F}M = \Lambda$ is a height one progenerator such that

$$0 \to \operatorname{Hom}_\Lambda(\mathbb{F}M,\mathbb{F}L) \to \operatorname{Hom}_\Lambda(\mathbb{F}M,\mathbb{F}M_0) \to \operatorname{Hom}_\Lambda(\mathbb{F}M,\mathbb{F}N) \to 0$$

is exact. By 4.4 we see that $\operatorname{End}_\Lambda(\mathbb{F}L \oplus \mathbb{F}M) \in \operatorname{CM} R$, hence $\operatorname{End}_R(L \oplus M) \cong \operatorname{End}_\Lambda(\mathbb{F}L \oplus \mathbb{F}M) \in \operatorname{CM} R$ as required. ☐

Now we are ready to prove the following crucial result (c.f. 5.13 later).

Proposition 4.6. *Let R be normal 3-CY$^-$ and let M be a non-zero modifying module. Then the following are equivalent*

(1) *M is a MM module.*

(2) *If N is any modifying module then there exists an exact sequence $0 \to M_1 \to M_0 \xrightarrow{f} N$ with each $M_i \in \operatorname{add} M$ such that f is a right $\operatorname{add} M$-approximation.*

Proof. Set $\Lambda := \mathrm{End}_R(M)$. Since M is a height one progenerator, we have a reflexive equivalence $\mathbb{F} := \mathrm{Hom}_R(M,-) : \mathrm{ref}\, R \to \mathrm{ref}\, \Lambda$ by 2.5(4). Moreover Λ is 3-CY$^-$ by 2.18 and so a Gorenstein R-order by 2.17.
(1)⇒(2) By taking a right $\mathrm{add}\, M$-approximation of N we have an exact sequence $0 \to L \to M_0 \xrightarrow{f} N$ with $M_0 \in \mathrm{add}\, M$ such that $0 \to \mathbb{F}L \to \mathbb{F}M_0 \xrightarrow{\mathbb{F}f} \mathbb{F}N \to 0$ is exact. By 4.5 $\mathrm{End}_R(L \oplus M) \in \mathrm{CM}\, R$ hence since M is a MM module, $L \in \mathrm{add}\, M$.
(2)⇒(1) Suppose N is reflexive with $\mathrm{End}_R(M \oplus N) \in \mathrm{CM}\, R$. Then $\mathbb{F}N \in \mathrm{CM}\, R$. We have $\mathrm{proj.dim}_\Lambda \mathbb{F}N \le 1$ since N is a modifying module and so there is an exact sequence $0 \to \mathbb{F}M_1 \to \mathbb{F}M_0 \to \mathbb{F}N \to 0$ by assumption. Since Λ is a Gorenstein R-order it follows that $\mathbb{F}N$ is a projective Λ-module by using localization and Auslander–Buchsbaum 2.13. Hence $N \in \mathrm{add}\, M$. ☐

The following version of the Bongartz completion is convenient for us. Recall from the introduction that throughout this paper when we say tilting module we mean a tilting module of projective dimension ≤ 1 (see 1.13).

Lemma 4.7. *Suppose R is normal, $M \in \mathrm{ref}\, R$ and denote $\Lambda := \mathrm{End}_R(M)$. If $N \in \mathrm{ref}\, R$ is such that $\mathrm{Hom}_R(M,N)$ is a partial tilting Λ-module then there exists $L \in \mathrm{ref}\, R$ such that $\mathrm{Hom}_R(M, N \oplus L)$ is a tilting Λ-module.*

Proof. By 2.5 $T := \mathrm{Hom}_R(M,N)$ and Λ are both reflexive. Thus since R is normal we can invoke [IR08, 2.8] to deduce that there exists an $X \in \mathrm{ref}\, \Lambda$ such that $T \oplus X$ is tilting. Again by 2.5 $X = \mathrm{Hom}_R(M,L)$ for some $L \in \mathrm{ref}\, R$. ☐

Proposition 4.8. *Let R be normal 3-CY$^-$ and suppose M is a MM module. Then*
(1) $\mathrm{Hom}_R(M,-)$ *sends modifying R-modules to partial tilting* $\mathrm{End}_R(M)$*-modules.*
(2) $\mathrm{Hom}_R(M,-)$ *sends MM R-modules to tilting* $\mathrm{End}_R(M)$*-modules.*

Proof. (1) Denote $\Lambda := \mathrm{End}_R(M)$, let N be a modifying module and denote $T := \mathrm{Hom}_R(M,N)$. Note first that $\mathrm{proj.dim}_\Lambda T \le 1$ by 4.6 and also Λ is a Gorenstein R-order by 2.17 and 2.18.

Since projective dimension localizes $\mathrm{proj.dim}_{\Lambda_\mathfrak{p}} T_\mathfrak{p} \le 1$ for all primes $\mathfrak{p}$ with $\mathrm{ht}\, \mathfrak{p} = 2$ and further for these primes $T_\mathfrak{p} \in \mathrm{CM}\, R_\mathfrak{p}$ by 2.3. Since $\Lambda_\mathfrak{p}$ is a Gorenstein $R_\mathfrak{p}$-order, Auslander–Buchsbaum (2.13) implies that $T_\mathfrak{p}$ is a projective $\Lambda_\mathfrak{p}$-module and so $\mathrm{Ext}^1_{\Lambda_\mathfrak{p}}(T_\mathfrak{p}, T_\mathfrak{p}) = 0$ for all primes $\mathfrak{p}$ with $\mathrm{ht}\, \mathfrak{p} = 2$. Consequently $\mathrm{Ext}^1_{\Lambda_\mathfrak{m}}(T_\mathfrak{m}, T_\mathfrak{m})$ has finite length for all $\mathfrak{m} \in \mathrm{Max}\, R$. On the other hand Λ is 3-CY$^-$ by 2.18 and $\mathrm{End}_\Lambda(T) \cong \mathrm{End}_R(N) \in \mathrm{CM}\, R$ by 2.5. Thus $\mathsf{fl}\, \mathrm{Ext}^1_{\Lambda_\mathfrak{m}}(T_\mathfrak{m}, T_\mathfrak{m}) = 0$ for all $\mathfrak{m} \in \mathrm{Max}\, R$ by 2.6 and so $\mathrm{Ext}^1_\Lambda(T,T) = 0$, as required.
(2) Now suppose that N is also MM. By Bongartz completion 4.7 we may find $L \in \mathrm{ref}\, R$ such that $\mathrm{Hom}_R(M, N \oplus L)$ is a tilting $\mathrm{End}_R(M)$-module, thus $\mathrm{End}_R(M)$ and $\mathrm{End}_R(N \oplus L)$ are derived equivalent. Since $\mathrm{End}_R(M)$ is 3-CY$^-$ so is $\mathrm{End}_R(N \oplus L)$ [IR08, 3.1(1)] and thus by 2.18 $\mathrm{End}_R(N \oplus L) \in \mathrm{CM}\, R$. Consequently $L \in \mathrm{add}\, N$ and so $\mathrm{Hom}_R(M,N)$ is a tilting module.

Now for the convenience of the reader we give a second proof, which shows us more explicitly how our tilting module generates the derived category. If N is also MM then since $(-)^* : \mathrm{ref}\, R \to \mathrm{ref}\, R$ is a duality, certainly N^* (and M^*) is MM. By 4.6 we can find

$$0 \to N_1^* \to N_0^* \to M^*$$

such that

$$0 \to \mathbb{F}N_1^* \to \mathbb{F}N_0^* \to \mathbb{F}M^* \to 0 \tag{4.D}$$

is exact, where $\mathbb{F} = \mathrm{Hom}_R(N^*,-)$. Denote $\Gamma := \mathrm{End}_R(N^*)$, then $\mathrm{Ext}^1_\Gamma(\mathbb{F}M^*, \mathbb{F}M^*) = 0$ by (1). Thus applying $\mathrm{Hom}_\Gamma(-, \mathbb{F}M^*)$ to (4.D) gives us the following commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{Hom}_\Gamma(\mathbb{F}M^*, \mathbb{F}M^*) \arrow[r] \arrow[d, equal] & \mathrm{Hom}_\Gamma(\mathbb{F}N_0^*, \mathbb{F}M^*) \arrow[r] \arrow[d, equal] & \mathrm{Hom}_\Gamma(\mathbb{F}N_1^*, \mathbb{F}M^*) \arrow[r] \arrow[d, equal] & 0 \\
0 \arrow[r] & \mathrm{Hom}_R(M^*, M^*) \arrow[r] & \mathrm{Hom}_R(N_0^*, M^*) \arrow[r] & \mathrm{Hom}_R(N_1^*, M^*) \arrow[r] & 0
\end{tikzcd}$$

where the top row is exact. Hence the bottom row is exact. Since $(-)^* : \operatorname{ref} R \to \operatorname{ref} R$ is a duality, this means that

$$0 \to \operatorname{Hom}_R(M, M) \to \operatorname{Hom}_R(M, N_0) \to \operatorname{Hom}_R(M, N_1) \to 0$$

is exact. But denoting $\Lambda := \operatorname{End}_R(M)$ and $T := \operatorname{Hom}_R(M, N)$, this means we have an exact sequence

$$0 \to \Lambda \to T_0 \to T_1 \to 0$$

with each $T_i \in \operatorname{add} T$. Hence T is a tilting Λ-module. $\square$

The following is now immediate:

Corollary 4.9. *Let R be normal 3-CY$^-$ and let M be a MM module. Then*
(1) *If N is any modifying module then the partial tilting $\operatorname{End}_R(M)$-module $T := \operatorname{Hom}_R(M, N)$ induces a recollement*

$$\begin{tikzcd}
\mathrm{U} \arrow[r] & \mathrm{D}(\operatorname{Mod}\operatorname{End}_R(M)) \arrow[l, bend right] \arrow[l, bend left] \arrow[r, "F"] & \mathrm{D}(\operatorname{Mod}\operatorname{End}_R(N)) \arrow[l, bend right] \arrow[l, bend left]
\end{tikzcd}$$

where $F = \mathbf{R}\operatorname{Hom}(T, -)$ and U is a certain triangulated subcategory of $\mathrm{D}(\operatorname{Mod}\operatorname{End}_R(M))$.
(2) *If further N is maximal modifying then the above functor F is an equivalence.*

Proof. (1) Set $\Lambda := \operatorname{End}_R(M)$ then $T := \operatorname{Hom}_R(M, N)$ is a partial tilting Λ-module by 4.8. The fact that $\operatorname{End}_\Lambda(T) \cong \operatorname{End}_R(N)$ follows since $\operatorname{Hom}_R(M, -)$ is a reflexive equivalence. By Bongartz completion T is a summand of a tilting Λ-module U. We have a derived equivalence $\mathrm{D}(\operatorname{Mod}\operatorname{End}_R(M)) \approx \mathrm{D}(\operatorname{Mod}\operatorname{End}_\Lambda(U))$. Moreover there exists an idempotent e of $\operatorname{End}_\Lambda(U)$ such that $e \operatorname{End}_\Lambda(U) e \cong \operatorname{End}_\Lambda(T) \cong \operatorname{End}_R(N)$. Thus we have the desired recollement (e.g. [M03, 2.17]).
(2) is an immediate consequence. $\square$

In the build-up to our next main results 4.14 and 4.15 we require four technical results (4.10, 4.11, 4.12 and 4.13) which ensure that membership of $\operatorname{add} M$ can be shown locally and on the derived level:

Lemma 4.10. *Let R be a commutative ring, $M, N \in \operatorname{mod} R$. Denote by $M^{\oplus m} \xrightarrow{f} N$ a right $\operatorname{add} M$-approximation of N, and by $N^{\oplus n} \xrightarrow{g} M$ a right $\operatorname{add} N$-approximation of M. Then $\operatorname{add} M \subseteq \operatorname{add} N$ if and only if the morphism $M^{\oplus mn} \xrightarrow{f^{\oplus n} g} M$ induces a surjection $\operatorname{Hom}_R(M, M^{\oplus mn}) \twoheadrightarrow \operatorname{End}_R(M)$.*

Proof. ($\Leftarrow$) If $\operatorname{Hom}_R(M, M^{\oplus mn}) \to \operatorname{End}_R(M)$ is surjective we may lift id_M to obtain a splitting for g and hence M is a summand of $N^{\oplus n}$.
($\Rightarrow$) If $M \in \operatorname{add} N$ then there exists $M \xrightarrow{a} N^{\oplus t} \xrightarrow{b} M$ with $ab = \mathrm{id}_M$. For every $\varphi \in \operatorname{End}_R(M)$ there is a commutative diagram

$$\begin{tikzcd}
M \arrow[r, "a"] \arrow[d, dotted, "\psi_1"] & N^{\oplus t} \arrow[r, "b"] \arrow[d, dotted, "\psi_0"] & M \arrow[d, "\varphi"] \\
M^{\oplus mn} \arrow[r, "f^{\oplus n}"] & N^{\oplus n} \arrow[r, "g"] & M
\end{tikzcd}$$

where ψ_0 exists since g is an approximation and ψ_1 exists since $f^{\oplus n}$ is an approximation. Consequently $\psi_1 f^{\oplus n} g = a\psi_0 g = ab\varphi = \varphi$ and so φ is the image of ψ_1 under the map $\operatorname{Hom}_R(M, M^{\oplus mn}) \to \operatorname{End}_R(M)$. Thus the map is surjective. $\square$

Proposition 4.11. *Let R be a commutative ring, $M, N \in \operatorname{mod} R$. Then the following are equivalent:*
1. $\operatorname{add} M \subseteq \operatorname{add} N$.
2. $\operatorname{add} M_{\mathfrak{p}} \subseteq \operatorname{add} N_{\mathfrak{p}}$ *for all* $\mathfrak{p} \in \operatorname{Spec} R$.
3. $\operatorname{add} M_{\mathfrak{m}} \subseteq \operatorname{add} N_{\mathfrak{m}}$ *for all* $\mathfrak{m} \in \operatorname{Max} R$.
4. $\operatorname{add} \widehat{M}_{\mathfrak{p}} \subseteq \operatorname{add} \widehat{N}_{\mathfrak{p}}$ *for all* $\mathfrak{p} \in \operatorname{Spec} R$.
5. $\operatorname{add} \widehat{M}_{\mathfrak{m}} \subseteq \operatorname{add} \widehat{N}_{\mathfrak{m}}$ *for all* $\mathfrak{m} \in \operatorname{Max} R$.

Furthermore we can replace $\subseteq$ by equality throughout and the result is still true.

Proof. Let f and g be as in 4.10. Then $f_{\mathfrak{p}} : M_{\mathfrak{p}}^{\oplus m} \to N_{\mathfrak{p}}$ is a right $\operatorname{add} M_{\mathfrak{p}}$-approximation and $\widehat{f_{\mathfrak{p}}} : \widehat{M_{\mathfrak{p}}}^{\oplus m} \to \widehat{N_{\mathfrak{p}}}$ is a right $\operatorname{add} \widehat{M_{\mathfrak{p}}}$-approximation for any $\mathfrak{p} \in \operatorname{Spec} R$. Similar statements hold for g. Since the vanishing of $\operatorname{Cok}(\operatorname{Hom}_R(M, M^{\oplus mn}) \xrightarrow{\cdot f^{\oplus n} g} \operatorname{End}_R(M))$ can be checked locally or complete locally, all conditions are equivalent. The last statement holds by symmetry. $\square$

Lemma 4.12. *Derived equivalences are preserved under localization and completion.*

Proof. Say A is derived equivalent to B via a tilting complex T, then since ext groups localize (respectively, complete), $T_{\mathfrak{p}}$ and $\widehat{T_{\mathfrak{p}}}$ both have no self-extensions. Further A can be reached from T using sums, summands and cones of T, so using the localizations (respectively, completions) of these triangles we conclude that $A_{\mathfrak{p}}$ can be reached from $T_{\mathfrak{p}}$ and also $\widehat{A_{\mathfrak{p}}}$ can be reached from $\widehat{T_{\mathfrak{p}}}$. Thus $T_{\mathfrak{p}}$ is a tilting $A_{\mathfrak{p}}$ complex and $\widehat{T_{\mathfrak{p}}}$ is a tilting $\widehat{A_{\mathfrak{p}}}$ complex. $\square$

Proposition 4.13. *Let Λ be a module finite R-algebra and let e be an idempotent of Λ. If $e\Lambda e$ is derived equivalent to Λ, then $\Lambda \in \operatorname{add} \Lambda e \subseteq \operatorname{mod} \Lambda$, i.e. Λe is a progenerator of Λ.*

Proof. By 4.11 we need to show that $\widehat{\Lambda}_{\mathfrak{p}} \in \operatorname{add} \widehat{\Lambda}_{\mathfrak{p}} e$ for all $\mathfrak{p} \in \operatorname{Spec} R$. There is a derived equivalence between $e\widehat{\Lambda}_{\mathfrak{p}}e$ and $\widehat{\Lambda}_{\mathfrak{p}}$ by 4.12. Since the derived equivalence gives an isomorphism between Grothendieck groups of perfect derived categories, the number of isomorphism classes of indecomposable projective $e\widehat{\Lambda}_{\mathfrak{p}}e$-modules is equal to that of $\widehat{\Lambda}_{\mathfrak{p}}$. Thus any indecomposable projective $\widehat{\Lambda}_{\mathfrak{p}}$-module appears in $\widehat{\Lambda}_{\mathfrak{p}}e$ as a summand, so we have $\widehat{\Lambda}_{\mathfrak{p}} \in \operatorname{add} \widehat{\Lambda}_{\mathfrak{p}}e$. $\square$

Corollary 4.14. *Let R be normal 3-CY$^-$ and let M be a MM module. Suppose $N \in \operatorname{ref} R$ such that $\operatorname{End}_R(M)$ is derived equivalent to $\operatorname{End}_R(N)$. Then N is also a MM module.*

Proof. By 2.18 $\operatorname{End}_R(M)$ is 3-CY$^-$ hence so is $\operatorname{End}_R(N)$ and thus $\operatorname{End}_R(N) \in \operatorname{CM} R$ again by 2.18. Hence N is a modifying module. Suppose that $X \in \operatorname{ref} R$ satisfies $\operatorname{End}_R(N \oplus X) \in \operatorname{CM} R$. Now 4.8(1) implies that $\operatorname{Hom}_R(M, N \oplus X)$ is a partial tilting $\operatorname{End}_R(M)$ module. By Bongartz completion 4.7 we may find $L \in \operatorname{ref} R$ such that $\operatorname{Hom}_R(M, N \oplus X \oplus L)$ is a tilting module and so $\operatorname{End}_R(N)$ and $\operatorname{End}_R(N \oplus X \oplus L) := \Lambda$ are derived equivalent since they are both derived equivalent to $\operatorname{End}_R(M)$. By 4.13

$$\operatorname{Hom}_R(N \oplus X \oplus L, N \oplus X \oplus L) = \Lambda \in \operatorname{add} \Lambda e = \operatorname{add} \operatorname{Hom}_R(N \oplus X \oplus L, N)$$

so since $\operatorname{Hom}(N \oplus X \oplus L, -) : \operatorname{ref} R \to \operatorname{ref} \Lambda$ is an equivalence, $N \oplus X \oplus L \in \operatorname{add} N$ and so in particular $X \in \operatorname{add} N$. $\square$

Theorem 4.15. *Let R be normal 3-CY$^-$ and let M be a MM module. Then the functor $\operatorname{Hom}_R(M, -) : \operatorname{mod} R \to \operatorname{mod} \operatorname{End}_R(M)$ induces bijections*

(1) $\{$*maximal modifying R-modules*$\} \overset{1:1}{\longleftrightarrow} \{$*reflexive tilting $\operatorname{End}_R(M)$-modules*$\}$.
(2) $\{$*modifying R-modules*$\} \overset{1:1}{\longleftrightarrow} \{$*reflexive partial tilting $\operatorname{End}_R(M)$-modules*$\}$.

Proof. (1) In light of 4.8(2) we only need to show that every reflexive tilting module is the image of some MM module. Thus let X be a reflexive tilting $\operatorname{End}_R(M)$-module; by reflexive equivalence there exists some $N \in \operatorname{ref} R$ such that $\operatorname{Hom}_R(M, N) \cong X$. We claim that N is MM. Since $\operatorname{Hom}_R(M, N)$ is a tilting $\operatorname{End}_R(M)$-module certainly $\operatorname{End}_R(M)$ and $\operatorname{End}_R(N)$ are derived equivalent; the fact that N is MM follows from 4.14 above.
(2) By 4.8(1) we only need to show that every reflexive partial tilting $\operatorname{End}_R(M)$ module is the image of some modifying module. Suppose X is a reflexive partial tilting $\operatorname{End}_R(M)$-module, say $X \cong \operatorname{Hom}_R(M, N)$. Then by Bongartz completion 4.7 there exists $N_1 \in \operatorname{ref} R$ such that $\operatorname{Hom}_R(M, N \oplus N_1)$ is a tilting $\operatorname{End}_R(M)$-module. By (1) $N \oplus N_1$ is MM, thus $\operatorname{End}_R(N)$ is a summand of the CM $\operatorname{End}_R(N \oplus N_1)$ and so is itself CM. $\square$

Corollary 4.16. *Let R be normal 3-CY$^-$ and let M be a MM module. Then*
(1) N is a modifying module $\iff$ N is the direct summand of a MM module.
(2) R has a MM module containing R as a summand. Necessarily this is CM.

Proof. (1) ‘if’ is clear. For the ‘only if’ let N be a modifying module, then by 4.8(1) $\mathrm{Hom}_R(M,N)$ is a partial tilting $\mathrm{End}_R(M)$-module, so the proof of 4.15(2) shows that there exists $N_1 \in \mathrm{ref}\, R$ such that $N \oplus N_1$ is MM.
(2) Apply (1) to R. The corresponding MM module is necessarily CM by 4.2. ☐

Recall from 1.5(3) we say that an R-order Λ has *isolated singularities* if $\mathrm{gl.dim}\, \Lambda_{\mathfrak{p}} = \dim R_{\mathfrak{p}}$ for all non-maximal primes $\mathfrak{p}$ of R.

Remark 4.17. It is unclear in what generality every maximal modification algebra $\mathrm{End}_R(M)$ has isolated singularities. In many cases this is true — for example if R is itself an isolated singularity this holds, as it does whenever M is CT by 5.5. Also, if R is Gorenstein, $X \to \mathrm{Spec}\, R$ with $M \in \mathrm{ref}\, R$ such that $\mathrm{D^b}(\mathrm{coh}\, X) \cong \mathrm{D^b}(\mathrm{mod}\, \mathrm{End}_R(M))$, then provided X has at worst isolated singularities (e.g. if X is a 3-fold with terminal singularities) then $\mathrm{End}_R(M)$ has isolated singularities too. This is a direct consequence of the fact that in this case the singular derived category has finite dimensional Hom-spaces. Also note that if R is normal 3-CY^- then the existence of a MM algebra $\mathrm{End}_R(M)$ with isolated singularities implies that $R_{\mathfrak{p}}$ has at worst ADE singularities for all primes $\mathfrak{p}$ of height 2 by a result of Auslander (see [IW08, 2.13]). Finally note that it follows immediately from 4.9 (and 4.12) that if R is normal 3-CY^- and there is one MM algebra $\mathrm{End}_R(M)$ with isolated singularities then necessary all MM algebras $\mathrm{End}_R(N)$ have isolated singularities.

The above remark suggests the following conjecture.

Conjecture 4.18. Let R be a normal 3-CY^- ring with rational singularities. Then
(1) R always has a MM module M (which may be R).
(2) For all such M, $\mathrm{End}_R(M)$ has isolated singularities.

This is closely related to a conjecture of Van den Bergh regarding the equivalence of the existence of crepant and noncommutative crepant resolutions when R is a rational normal Gorenstein 3-fold. We remark that given the assumption on rational singularities, any proof is likely to be geometric.

5. Relationship between CT modules, NCCRs and MM modules

In this section we define CT modules for non-isolated singularities and we show that they are a special case of the MM modules introduced in §4. In §5.2 we show that all these notions recover the established ones when R is an isolated singularity.

When R is a normal 3-CY^- domain, below we prove the implications in the following picture which summarizes the relationship between CT modules, NCCRs and MM modules:

$$\begin{tikzcd}
\text{CT modules} \arrow[r, Rightarrow, shift left, "5.5"] & \text{modules giving NCCRs} \arrow[l, Rightarrow, shift left, "\substack{\text{if generator}\\ 5.5}"] \arrow[r, Rightarrow, shift left, "5.1"] & \text{MM modules} \arrow[l, Rightarrow, shift left, "\substack{\text{if }\exists\text{NCCR}\\ 5.14}"] \arrow[r, Rightarrow] & \text{modifying modules}
\end{tikzcd}$$

First we have the following, which is true in arbitrary dimension:

Proposition 5.1. *Let R be a d-dimensional, normal, equi-codimensional CM ring (e.g. if R is normal d-CY^-) with a canonical module ω_R. Then reflexive R-modules M giving NCCRs are MM modules.*

Proof. Assume that $X \in \mathrm{ref}\, R$ satisfies $\mathrm{End}_R(M \oplus X) \in \mathrm{CM}\, R$. Then $\mathrm{Hom}_R(M,X) \in \mathrm{CM}\, \Gamma$ for $\Gamma := \mathrm{End}_R(M)$. By 2.14 we have $\mathrm{Hom}_R(M,X) \in \mathrm{proj}\, \Gamma$. By 2.5(4) $X \in \mathrm{add}\, M$ as required. ☐

5.1. CT Modules on Non-Isolated Singularities.

Definition 5.2. *Let R be a d-dimensional CM ring with a canonical module ω_R. We call $M \in \mathrm{CM}\, R$ a CT module if*

$$\mathrm{add}\, M = \{X \in \mathrm{CM}\, R : \mathrm{Hom}_R(M,X) \in \mathrm{CM}\, R\} = \{X \in \mathrm{CM}\, R : \mathrm{Hom}_R(X,M) \in \mathrm{CM}\, R\}.$$

We have the following easy observations.

Lemma 5.3. (1) *Any CT module is a generator-cogenerator.*
(2) *Any CT module is maximal modifying.*

Proof. Let M be a CT module.
(1) This is clear from $\mathrm{Hom}_R(M,\omega_R)\in \mathrm{CM}\,R$ and $\mathrm{Hom}_R(R,M)\in\mathrm{CM}\,R$.
(2) Suppose $N\in\mathrm{ref}\,R$ with $\mathrm{End}_R(M\oplus N)\in\mathrm{CM}\,R$, then certainly $\mathrm{Hom}_R(M,N)\in\mathrm{CM}\,R$. Since $R\in\mathrm{add}\,M$ by (1), we have $N\in\mathrm{CM}\,R$. Hence since M is a CT module, necessarily $N\in\mathrm{add}\,M$. $\square$

Not every MM module is CT, however in the case when R has a CT module (which implies R has a NCCR by 5.12(3) below, which in turn implies by [V04b] that R has a crepant resolution) we give a rather remarkable relationship between CT modules, MM modules and NCCRs in 5.14 at the end of this subsection.

If R is a CM ring with a canonical module ω_R we denote the duality $(-)^\vee:=\mathrm{Hom}_R(-,\omega_R)$. We shall see shortly that if M or $M^\vee$ is a generator then we may test the above CT condition on one side (see 5.5 below), but before we do this we need the following easy observation.

Lemma 5.4. *Let R be a CM ring with a canonical module ω_R, $M\in\mathrm{CM}\,R$. If $\mathrm{End}_R(M)$ is a non-singular R-order, then $R\in\mathrm{add}\,M\iff\omega_R\in\mathrm{add}\,M$.*

Proof. Since $(-)^\vee:\mathrm{CM}\,R\to\mathrm{CM}\,R$ is a duality we know that $\mathrm{End}_R(M^\vee)\cong\mathrm{End}_R(M)^{\mathrm{op}}$ and so $\mathrm{End}_R(M^\vee)$ is also a non-singular R-order. Moreover $R^\vee=\omega_R$ and $\omega_R^\vee=R$ so by the symmetry of this situation we need only prove the 'only if' part. Thus assume that $R\in\mathrm{add}\,M$. In this case since $\mathrm{Hom}_R(M,\omega_R)=M^\vee\in\mathrm{CM}\,R$, by 2.14 $\mathrm{Hom}_R(M,\omega_R)$ is a projective $\mathrm{End}_R(M)$-module and thus $\omega_R\in\mathrm{add}\,M$ by 2.5(1). $\square$

We reach one of our main characterizations of CT modules. Note that if R is 3-CY$^-$ then by 2.8 the left-right symmetry in the definition of CT modules is immediate, however the following argument works in a little more generality:

Theorem 5.5. *Let R be a d-dimensional, equi-codimensional CM ring (e.g. if R is d-CY$^-$) with a canonical module ω_R. Then for any $M\in\mathrm{CM}\,R$ the following are equivalent*
(1) *M is a CT module.*
(2) *$R\in\mathrm{add}\,M$ and $\mathrm{add}\,M=\{X\in\mathrm{CM}\,R:\mathrm{Hom}_R(M,X)\in\mathrm{CM}\,R\}$.*
(2)$'$ *$\omega_R\in\mathrm{add}\,M$ and $\mathrm{add}\,M=\{X\in\mathrm{CM}\,R:\mathrm{Hom}_R(X,M)\in\mathrm{CM}\,R\}$.*
(3) *$R\in\mathrm{add}\,M$ and $\mathrm{End}_R(M)$ is a non-singular R-order.*
(3)$'$ *$\omega_R\in\mathrm{add}\,M$ and $\mathrm{End}_R(M)$ is a non-singular R-order.*
In particular CT modules are precisely the CM generators which give NCCRs.

Proof. (2)⇒(3). $R\in\mathrm{add}\,M$ by assumption. By (2) necessarily $\mathrm{End}_R(M)\in\mathrm{CM}\,R$. Now let $Y\in\mathrm{mod}\,\mathrm{End}_R(M)$ and consider a projective resolution $P_{d-1}\to P_{d-2}\to\ldots\to P_0\to Y\to 0$. By 2.5(1) there is an exact sequence $M_{d-1}\xrightarrow{f}M_{d-2}\to\ldots\to M_0$ with each $M_i\in\mathrm{add}\,M$ such that the projective resolution above is precisely

$$\mathrm{Hom}_R(M,M_{d-1})\xrightarrow{\cdot f}\mathrm{Hom}_R(M,M_{d-2})\to\ldots\to\mathrm{Hom}_R(M,M_0)\to Y\to 0.$$

Denote $K_d=\mathrm{Ker}\,f$. Then we have an exact sequence

$$0\to\mathrm{Hom}_R(M,K_d)\to P_{d-1}\to P_{d-2}\to\ldots\to P_0\to Y\to 0.$$

Localizing the above and counting depths we see that $\mathrm{Hom}_R(M,K_d)_{\mathfrak{m}}\in\mathrm{CM}\,R_{\mathfrak{m}}$ for all $\mathfrak{m}\in\mathrm{Max}\,R$, thus $\mathrm{Hom}_R(M,K_d)\in\mathrm{CM}\,R$ and so by definition $K_d\in\mathrm{add}\,M$. Hence $\mathrm{proj.dim}_{\mathrm{End}_R(M)}Y\le d$ and so $\mathrm{gl.dim}\,\mathrm{End}_R(M)\le d$.
(3)⇒(2). Since $\mathrm{End}_R(M)\in\mathrm{CM}\,R$, automatically $\mathrm{add}\,M\subseteq\{X\in\mathrm{CM}\,R:\mathrm{Hom}_R(M,X)\in\mathrm{CM}\,R\}$. To obtain the reverse inclusion assume that $X\in\mathrm{CM}\,R$ with $\mathrm{Hom}_R(M,X)\in\mathrm{CM}\,R$, then since $\mathrm{End}_R(M)$ is a non-singular R-order $\mathrm{Hom}_R(M,X)$ is a projective $\mathrm{End}_R(M)$ module by 2.14. This implies that $X\in\mathrm{add}\,M$ by 2.5(1).
(2)$'$ $\iff$ (3)$'$ We have a duality $(-)^\vee:\mathrm{CM}\,R\to\mathrm{CM}\,R$ thus apply (2) $\iff$ (3) to $M^\vee$ and use the fact that $\mathrm{End}_R(M^\vee)=\mathrm{End}_R(M)^{\mathrm{op}}$ has finite global dimension if and only if $\mathrm{End}_R(M)$ does.
(3) $\iff$ (3)$'$ This is immediate from 5.4.

In particular by the above we have (2) $\iff$ (2)$'$. Since we clearly have (1) $\iff$ (2)+(2)$'$, the proof is completed. □

Note that the last assertion in 5.5 is improved when R is a 3-CY$^-$ normal domain in 5.14(3). From the definition it is not entirely clear that CT is a local property:

Corollary 5.6. *Let R be a d-dimensional, equi-codimensional CM ring (e.g. if R is d-CY$^-$) with a canonical module ω_R Then M is a CT R-module if and only if $M_{\mathfrak{p}}$ is a CT $R_{\mathfrak{p}}$-module for all primes $\mathfrak{p}$. Thus CT is a local property.*

Proof. By 5.5 M is a CT R-module if and only if $R \in \operatorname{add} M$ and $\operatorname{End}_R(M)$ is a non-singular R-order. By 4.11 this is equivalent to that $R_{\mathfrak{p}} \in \operatorname{add} M_{\mathfrak{p}}$ and $\operatorname{End}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) = \operatorname{End}_R(M)_{\mathfrak{p}}$ is a non-singular $R_{\mathfrak{p}}$-order for all $\mathfrak{p} \in \operatorname{Spec} R$. Again by 5.5 this is equivalent to that $M_{\mathfrak{p}}$ is a CT $R_{\mathfrak{p}}$-module for all $\mathfrak{p} \in \operatorname{Spec} R$. □

Theorem 5.5 also gives an easy method to find examples of CT modules. Recall that an element $g \in \operatorname{GL}(d,k)$ is called *pseudo-reflection* if the rank of $g-1$ is at most one. A finite subgroup G of $\operatorname{GL}(d,k)$ is called *small* if it does not contain pseudo-reflections except the identity. The following is well-known:

Proposition 5.7. *Let k be a field of characteristic zero, let S be the polynomial ring $k[x_1,\ldots,x_d]$ (resp. the formal power series ring $k[[x_1,\ldots,x_d]]$) and let G be a finite subgroup of $\operatorname{GL}(d,k)$.*
(1) If G is generated by pseudo-reflections, then S^G is a polynomial ring (resp. a formal power series ring) in d variables.
(2) If G is small, then the natural map $S\#G \to \operatorname{End}_R(S)$ given by $sg \mapsto (t \mapsto s\cdot g(t))$ $(s,t \in S,\ g \in G)$ is an isomorphism.

Proof. (1) See [Bou68] for example.
(2) This is due to Auslander [Aus86, Yos90] (see [IT] for a detailed proof). □

This immediately gives us a rich source of CT modules:

Theorem 5.8. *Let k be a field of characteristic zero, and let S be the polynomial ring $k[x_1,\ldots,x_d]$ (resp. the formal power series ring $k[[x_1,\ldots,x_d]]$). For a finite subgroup G of $\operatorname{GL}(d,k)$, let $R = S^G$. If G is small, then S is a CT R-module.*

Proof. Since G is small $\operatorname{End}_R(S) \cong S\#G$ by 5.7(2). Thus by 2.10 $\operatorname{End}_R(S)$ is a non-singular R-order and so consequently S is a CT R-module by 5.5. □

We can say more in the complete local setting. The following result is shown in [Iya07, 2.5] under the assumption that G is a small subgroup of $\operatorname{GL}(d,k)$ and S^G is an isolated singularity. We can drop both assumptions under our definition of CT modules.

Theorem 5.9. *Let k be a field of characteristic zero and let $S := k[[x_1,\ldots,x_d]]$ be the formal power series ring. For an arbitrary finite group G together with a group homomorphism $G \to \operatorname{Aut}_{k-\mathrm{alg}}(S)$, let $R := S^G$. Then S is a CT R-module.*

In the proof, we need the following easy lemma.

Lemma 5.10. *Let k be a field, and let $S := k[[x_1,\ldots,x_d]]$ be the formal power series ring. For an arbitrary finite group G together with a group homomorphism $G \to \operatorname{Aut}_{k-\mathrm{alg}}(S)$ such that $|G| \neq 0$ in k, there exist elements $y_1,\ldots,y_d \in S$ such that S is the formal power series ring $k[[y_1,\ldots,y_d]]$ with variables $y_1,\ldots,y_d$ and G acts linearly on the subspace $\sum_{i=1}^d ky_i$ of S.*

Proof. For the convenience of the reader, we include a proof.

We denote the maximal ideal of S by $\mathfrak{n}$. Let $V := \mathfrak{n}/\mathfrak{n}^2$. This is a k-vector space with a basis $x_1,\ldots,x_d$. We naturally regard V as a subspace of S. Clearly we have $g\mathfrak{n} = \mathfrak{n}$ and $g\mathfrak{n}^2 = \mathfrak{n}^2$ for any $g \in G$. Thus we have an induced action of G on the k-vector space V. For each $g \in G$, we denote by $g' \in \operatorname{Aut}_k(V)$ the induced action.

For all $g \in G$ and all i with $1 \le i \le d$, we have

$$g'x_i - gx_i \in \mathfrak{n}^2,$$

so we have

$$g^{-1}g'x_i - x_i \in g^{-1}\mathfrak{n}^2 = \mathfrak{n}^2.$$

For all i with $1 \le i \le d$ we define

$$y_i := \frac{1}{|G|}\sum_{g\in G} g^{-1}(g'x_i),$$

then we have $y_i - x_i \in \mathfrak{n}^2$. Hence S is the formal power series ring $k[[y_1,\ldots,y_d]]$ with variables $y_1,\ldots,y_d$.

Since $(-)' : G \to \mathrm{Aut}_k(V)$ is a group homomorphism, we have $g' = (gh^{-1})'h'$ for all $h, g \in G$. Thus for all $h \in G$ and all i with $1 \le i \le d$,

$$hy_i = \frac{1}{|G|}\sum_{g\in G} hg^{-1}(g'x_i) = \frac{1}{|G|}\sum_{g\in G}(gh^{-1})^{-1}((gh^{-1})'(h'x_i)) = \frac{1}{|G|}\sum_{g\in G} g^{-1}(g'(h'x_i)).$$

Thus $h'x_i \in V = \sum_{i=1}^{d} kx_i$ implies that $hy_i \in \sum_{i=1}^{d} ky_i$, as required. □

Now we are ready to prove 5.9. We use induction on $|G|$. Denote the kernel of $G \to \mathrm{Aut}_{k-\mathrm{alg}}(S)$ by K, then since $R = S^G = S^{G/K}$ we can assume that the action is faithful. By 5.10, we can assume that G is a subgroup of $\mathrm{GL}(d,k)$.

If G is small, then we have $\mathrm{End}_R(S) \simeq S\#G$ by 5.7(2). Since $S\#G$ is a non-singular R-order by 2.10, we have the assertion by 5.5.

Assume that G is not small. Let H be the subgroup of G generated by pseudo-reflections and let $T := S^H$. Then H is a normal subgroup of G and there exists a group homomorphism $G/H \to \mathrm{Aut}_{k-\mathrm{alg}}(T)$ such that $R = T^{G/H}$. By 5.7(1), T is isomorphic to a formal power series ring of d variables. Thus S is a free T-module of finite rank. By induction T is a CT R-module. Thus S is also a CT R-module, since it is a direct sum of copies of T. □

As another source of CT modules, we have:

Example 5.11. Let $Y \xrightarrow{f} X = \mathrm{Spec}\, R$ be a projective birational morphism such that $Rf_*\mathscr{O}_Y = \mathscr{O}_X$ and every fibre has dimension ≤ 1, where R is a 3-dimensional normal Gorenstein affine ring R . Then provided Y is smooth and crepant there exists a NCCR $\mathrm{End}_R(M)$ [V04a, 3.2.10] in which M is CM containing R as a summand. By 5.5 M is a CT module.

We now show that for R normal 3-CY$^-$, the existence of a CT module is equivalent to the existence of a NCCR. Note that (2) below answers a question of Van den Bergh [V04b].

Corollary 5.12. *Let R be a 3-CY$^-$ normal domain. Then*
(1) CT modules are precisely those reflexive generators which give NCCRs.
(2) R has a NCCR $\iff$ R has a NCCR given by a CM generator M $\iff$ $\mathrm{CM}\, R$ contains a CT module.

Proof. Notice that any reflexive generator M which gives a NCCR is CM since R is a summand of M and further $M \cong \mathrm{Hom}_R(R,M)$ is a summand of $\mathrm{End}_R(M) \in \mathrm{CM}\, R$ as an R-module.
(1) By 5.5 CT modules are precisely the CM generators which give NCCRs. The assertion follows from the above remark.
(2) The latter equivalence was shown in 5.5. We only have to show ($\Rightarrow$) of the former assertion. By [IR08, 8.9(4)] if R has a NCCR then there exists a reflexive generator which give a NCCR. This is CM by the above remark. □

Below is another characterization of CT modules. Compare this to the previous 4.6.

Proposition 5.13. *Assume R is a 3-CY$^-$ normal domain and let $M \in \mathrm{CM}\, R$ with $R \in \mathrm{add}\, M$. Then the following are equivalent*
(1) M is a CT module.
(2) $\mathrm{End}_R(M) \in \mathrm{CM}\, R$ and further for all $X \in \mathrm{CM}\, R$ there exists an exact sequence $0 \to M_1 \to M_0 \xrightarrow{f} X \to 0$ with $M_1, M_0 \in \mathrm{add}\, M$ and a right $\mathrm{add}\, M$-approximation f.

Proof. (1)⇒(2). Fix $X \in \operatorname{CM} R$. Since R is 3-CY$^-$, we have an exact sequence $0 \to X \to P_0 \to P_1$ with each $P_i \in \operatorname{add} R$. Applying $\operatorname{Hom}_R(M,-)$ to this sequence shows that $\operatorname{proj.dim}_{\operatorname{End}_R(M)} \operatorname{Hom}_R(M,X) \leq 1$ since we know that $\operatorname{gl.dim} \operatorname{End}_R(M) = 3$ by 5.5. Consequently we may take a projective resolution $0 \to \operatorname{Hom}_R(M,M_1) \to \operatorname{Hom}_R(M,M_0) \to \operatorname{Hom}_R(M,X) \to 0$ which necessarily comes from a complex $0 \to M_1 \to M_0 \to X \to 0$. This complex is itself exact since M is a generator.
(2)⇒(1). Denote $\Gamma = \operatorname{End}_R(M)$. By 2.18 and 2.17 Γ is a Gorenstein R-order. By 5.5 we only have to show that $\operatorname{add} M = \{X \in \operatorname{CM} R : \operatorname{Hom}_R(M,X) \in \operatorname{CM} R\}$. The assumption $\operatorname{End}_R(M) \in \operatorname{CM} R$ shows that the inclusion $\subseteq$ holds so let $X \in \operatorname{CM} R$ be such that $\operatorname{Hom}_R(M,X) \in \operatorname{CM} R$. By assumption we may find $M_1, M_0 \in \operatorname{add} M$ such that $0 \to \operatorname{Hom}_R(M,M_1) \to \operatorname{Hom}_R(M,M_0) \to \operatorname{Hom}_R(M,X) \to 0$ is exact, hence $\operatorname{proj.dim}_{\Gamma_{\mathfrak{m}}} \operatorname{Hom}_R(M,X)_{\mathfrak{m}} < \infty$ for all $\mathfrak{m} \in \operatorname{Max} R$. Since $\operatorname{Hom}_R(M,X)_{\mathfrak{m}} \in \operatorname{CM} R_{\mathfrak{m}}$, Auslander–Buchsbaum (2.13) implies that $\operatorname{proj.dim}_{\Gamma_{\mathfrak{m}}} \operatorname{Hom}_R(M,X)_{\mathfrak{m}} = 0$ for all $\mathfrak{m} \in \operatorname{Max} R$ and hence $\operatorname{Hom}_R(M,X)$ is a projective Γ-module. Since M is a generator, $X \in \operatorname{add} M$. □

Provided an NCCR exists, the following shows the precise relationship between MM modules, CT modules and NCCRs. Note that 5.14(2) says that CT modules are really a special case of MM modules.

Proposition 5.14. *Let R be a 3-CY$^-$ normal domain, and assume that R has a NCCR (equivalently, by 5.12, a CT module). Then*
(1) *MM modules are precisely the reflexive modules which give NCCRs.*
(2) *MM modules which are CM (equivalently, by 4.2, the MM generators) are precisely the CT modules.*
(3) *CT modules are precisely those CM modules which give NCCRs.*

Proof. (1) (⇐) This is shown in 5.1 above.
(⇒) If M is a MM module then $\operatorname{End}_R(M) \in \operatorname{CM} R$ and so by [IR08, 8.9(3)] M is the summand of some reflexive module giving an NCCR. Thus $\operatorname{End}_R(M \oplus L)$ is a NCCR for some reflexive module L, so in particular $\operatorname{End}_R(M \oplus L) \in \operatorname{CM} R$. This implies that $L \in \operatorname{add} M$ and so in fact $\operatorname{End}_R(M)$ gives an NCCR.
(2) By (1) MM generators are precisely the CM generators which give NCCRs. By 5.5 these are precisely the CT modules.
(3) Follows immediately from (1) and (2). □

5.2. **Modifying Modules on Isolated Singularities.** In this subsection we relate our work to that of the more common notions of rigid, maximal rigid and maximal 1-orthogonal modules, and show that when R is an isolated singularity our new definitions of modifying, maximal modifying and CT modules recover the old ones.

Recall that $M \in \operatorname{ref} R$ is called *rigid* if $\operatorname{Ext}^1_R(M,M) = 0$. We call $M \in \operatorname{ref} R$ *maximal rigid* if M is rigid and furthermore it is maximal with respect to this property, namely if there exists $X \in \operatorname{ref} R$ such that $M \oplus X$ is rigid, then $X \in \operatorname{add} M$.

Recall that $M \in \operatorname{CM} R$ is called a *maximal* 1*-orthogonal* module if

$$\operatorname{add} M = \{X \in \operatorname{CM} R : \operatorname{Ext}^1_R(M,X) = 0\} = \{X \in \operatorname{CM} R : \operatorname{Ext}^1_R(X,M) = 0\}.$$

The key observation is the following.

Lemma 5.15. *Let R be a CM ring with only isolated singularities. Then $\operatorname{Ext}^1_R(X,Y)$ is a finite length R-module for any $X, Y \in \operatorname{CM} R$.*

Proof. This is well-known [Aus78, Yos90]. □

Proposition 5.16. *Let R be 3-CY$^-$ with only isolated singularities, $M \in \operatorname{CM} R$. Then*
(1) *M is a modifying module if and only if it is rigid.*
(2) *M is a maximal modifying module if and only if it is maximal rigid.*
(3) *M is a CT module if and only if it is maximal* 1*-orthogonal.*

Proof. Let $X, Y \in \operatorname{CM} R$. By 2.6 and 5.15 $\operatorname{Hom}_R(X,Y) \in \operatorname{CM} R$ if and only if $\operatorname{fl} \operatorname{Ext}^1_R(X,Y) = 0$ if and only if $\operatorname{Ext}^1_R(X,Y) = 0$. Thus the assertions follow. □

6. Mutations of Modifying Modules

6.1. **General Definitions and First Properties.** Mutation is a technique used to obtain new modifying, maximal modifying and CT modules from a given one. Many of our arguments work in the full generality of modifying modules athough sometimes it is necessary to restrict to the maximal modifying level to apply certain arguments. Throughout this section R will be a normal 3-CY$^-$ ring, M will be a modifying module with N such that $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$. Note that N may or may not be decomposable. Given this, we define left and right mutation as in 1.19 in the introduction; we have exact sequences

$$0 \to K_0 \xrightarrow{c} N_0 \xrightarrow{a} M \tag{6.E}$$

$$0 \to K_1 \xrightarrow{d} N_1^* \xrightarrow{b} M^* \tag{6.F}$$

where a is a right $\operatorname{add} N$-approximation and b is a right $\operatorname{add} N^*$-approximation. Thus $N_0, N_1 \in \operatorname{add} N$ and we have exact sequences

$$0 \to \operatorname{Hom}_R(N, K_0) \xrightarrow{\cdot c} \operatorname{Hom}_R(N, N_0) \xrightarrow{\cdot a} \operatorname{Hom}_R(N, M) \to 0 \tag{6.G}$$

$$0 \to \operatorname{Hom}_R(N^*, K_1) \xrightarrow{\cdot d} \operatorname{Hom}_R(N^*, N_1^*) \xrightarrow{\cdot b} \operatorname{Hom}_R(N^*, M^*) \to 0 \tag{6.H}$$

From this we define $\mu_N(M) := N \oplus K_0$ and $\nu_N(M) := N \oplus K_1^*$.

In general right and left mutation are not the same, although in some cases we will see that $\mu_N = \nu_N$ (6.14). Note it is certainly possible that $\mu_N(M) = \nu_N(M) = M$ and so the mutation is trivial; a concrete example is given by taking $R = k[x,y,z]^G$ with $G = \frac{1}{2}(1,1,0)$, $M = k[x,y,z]$ and $N = R$.

Firstly, we note that mutation is unique up to additive closure. This can be improved when R is complete local.

Lemma 6.1. *Suppose $N_0 \xrightarrow{a} M$ and $N_0' \xrightarrow{a'} M$ are two right $\operatorname{add} N$-approximations of M. Then $\operatorname{add}(N \oplus \operatorname{Ker} a) = \operatorname{add}(N \oplus \operatorname{Ker} a')$. A similar statement holds for left approximations.*

Proof. Let $K := \operatorname{Ker} a$ and $K' := \operatorname{Ker} a'$. Then we have a commutative diagram

$$\begin{tikzcd} 0 \arrow[r] & K \arrow[r, "c"] \arrow[d, "s"] & N_0 \arrow[r, "a"] \arrow[d, "t"] & M \arrow[d, equal] \\ 0 \arrow[r] & K' \arrow[r, "c'"] & N_0' \arrow[r, "a'"] & M \end{tikzcd}$$

of exact sequences, giving an exact sequence

$$0 \to K \xrightarrow{(-s\ c)} K' \oplus N_0 \xrightarrow{\binom{c'}{t}} N_0'. \tag{6.I}$$

From the commutative diagram

$$\begin{tikzcd} 0 \arrow[r] & \operatorname{Hom}_R(N,K) \arrow[r, "\cdot c"] \arrow[d, "\cdot s"] & \operatorname{Hom}_R(N,N_0) \arrow[r, "\cdot a"] \arrow[d, "\cdot t"] & \operatorname{Hom}_R(N,M) \arrow[r] \arrow[d, equal] & 0 \\ 0 \arrow[r] & \operatorname{Hom}_R(N,K') \arrow[r, "\cdot c'"] & \operatorname{Hom}_R(N,N_0') \arrow[r, "\cdot a'"] & \operatorname{Hom}_R(N,M) \arrow[r] & 0 \end{tikzcd}$$

we see that

$$\operatorname{Hom}_R(N, K' \oplus N_0) \xrightarrow{\cdot\binom{c'}{t}} \operatorname{Hom}_R(N, N_0') \to 0$$

is exact. Thus (6.I) is a split short exact sequence, so in particular $K \in \operatorname{add}(N \oplus K')$. Similarly $K' \in \operatorname{add}(N \oplus K)$. □

Proposition 6.2. *Let R be normal 3-CY$^-$ and let M be a modifying module with N such that $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$. Then both $\mu_N(M)$ and $\nu_N(M)$ are modifying R-modules.*

Proof. Immediate from 4.5. □

Proposition 6.3. *Let R be normal 3-CY$^-$ and let M be a modifying module with $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$ (i.e. notation as above). Then*

(1) *Applying $\operatorname{Hom}_R(-, N)$ to (6.E) induces an exact sequence*

$$0 \to \operatorname{Hom}_R(M, N) \xrightarrow{a\cdot} \operatorname{Hom}_R(N_0, N) \xrightarrow{c\cdot} \operatorname{Hom}_R(K_0, N) \to 0. \tag{6.J}$$

In particular c is a left $\operatorname{add} N$*-approximation.*
(2) *Applying* $\operatorname{Hom}_R(-,N^*)$ *to* (6.F) *induces an exact sequence*

$$0\to \operatorname{Hom}_R(M^*,N^*)\xrightarrow{b\cdot}\operatorname{Hom}_R(N_1^*,N^*)\xrightarrow{d\cdot}\operatorname{Hom}_R(K_1,N^*)\to 0 \tag{6.K}$$

In particular d is a left $\operatorname{add} N^*$*-approximation.*
(3) *We have that*

$$0\to M^*\xrightarrow{a^*}N_0^*\xrightarrow{c^*}K_0^* \tag{6.L}$$

$$0\to M\xrightarrow{b^*}N_1\xrightarrow{d^*}K_1^* \tag{6.M}$$

are exact, inducing exact sequences

$$0\to \operatorname{Hom}_R(N^*,M^*)\xrightarrow{\cdot a^*}\operatorname{Hom}_R(N^*,N_0^*)\xrightarrow{\cdot c^*}\operatorname{Hom}_R(N^*,K_0^*)\to 0 \tag{6.N}$$

$$0\to \operatorname{Hom}_R(K_0^*,N^*)\xrightarrow{c^*\cdot}\operatorname{Hom}_R(N_0^*,N^*)\xrightarrow{a^*\cdot}\operatorname{Hom}_R(M^*,N^*)\to 0 \tag{6.O}$$

$$0\to \operatorname{Hom}_R(N,M)\xrightarrow{\cdot b^*}\operatorname{Hom}_R(N,N_1)\xrightarrow{\cdot d^*}\operatorname{Hom}_R(N,K_1^*)\to 0 \tag{6.P}$$

$$0\to \operatorname{Hom}_R(K_1^*,N)\xrightarrow{d^*\cdot}\operatorname{Hom}_R(N_1,N)\xrightarrow{b^*\cdot}\operatorname{Hom}_R(M,N)\to 0 \tag{6.Q}$$

Proof. (1) Denote $\Lambda:=\operatorname{End}_R(N)$ and $\mathbb{F}:=\operatorname{Hom}_R(N,-)$. Then (6.G) is

$$0\to\mathbb{F}K_0\to\mathbb{F}N_0\to\mathbb{F}M\to 0$$

so applying $\operatorname{Hom}_\Lambda(-,\mathbb{F}N)$ gives

$$0\to\operatorname{Hom}_\Lambda(\mathbb{F}M,\mathbb{F}N)\to\operatorname{Hom}_\Lambda(\mathbb{F}N_0,\mathbb{F}N)\to\operatorname{Hom}_\Lambda(\mathbb{F}K_0,\mathbb{F}N)\to\operatorname{Ext}^1_\Lambda(\mathbb{F}M,\Lambda).$$

But by 2.18 Λ is 3-CY$^-$ and thus a Gorenstein R-order. Since $\mathbb{F}M\in\operatorname{CM}\Lambda$ it follows that $\operatorname{Ext}^1_\Lambda(\mathbb{F}M,\Lambda)=0$ and hence we have a commutative diagram of complexes

$$\begin{tikzcd}
0 \arrow[r] & \operatorname{Hom}_\Lambda(\mathbb{F}M,\mathbb{F}N) \arrow[r] \arrow[d, equal] & \operatorname{Hom}_\Lambda(\mathbb{F}N_0,\mathbb{F}N) \arrow[r] \arrow[d, equal] & \operatorname{Hom}_\Lambda(\mathbb{F}K_0,\mathbb{F}N) \arrow[r] \arrow[d, equal] & 0 \\
0 \arrow[r] & \operatorname{Hom}_R(M,N) \arrow[r, "a\cdot"] & \operatorname{Hom}_R(N_0,N) \arrow[r, "c\cdot"] & \operatorname{Hom}_R(K_0,N) \arrow[r] & 0
\end{tikzcd}$$

in which the top row is exact. It follows that the bottom row is exact.
(2) is identical to (1) since $\operatorname{Hom}_R(N^*,M^*)\in\operatorname{CM}R$.
(3) As in (1) denote $\Lambda:=\operatorname{End}_R(N)$ and $\mathbb{F}:=\operatorname{Hom}_R(N,-)$, then applying $\operatorname{Hom}_\Lambda(-,\mathbb{F}R)$ to (6.G) gives an commutative diagram of complexes

$$\begin{tikzcd}
0 \arrow[r] & \operatorname{Hom}_\Lambda(\mathbb{F}M,\mathbb{F}R) \arrow[r] \arrow[d, equal] & \operatorname{Hom}_\Lambda(\mathbb{F}N_0,\mathbb{F}R) \arrow[r] \arrow[d, equal] & \operatorname{Hom}_\Lambda(\mathbb{F}K_0,\mathbb{F}R) \arrow[d, equal] \\
0 \arrow[r] & \operatorname{Hom}_R(M,R) \arrow[r, "a^*"] & \operatorname{Hom}_R(N_0,R) \arrow[r, "c^*"] & \operatorname{Hom}_R(K_0,R)
\end{tikzcd}$$

in which the top row is exact. Hence the bottom row (i.e. (6.L)) is exact. The proof that (6.M) is exact is identical. Now since $(-)^*:\operatorname{ref}R\to\operatorname{ref}R$ is a duality, the sequences (6.N), (6.O), (6.P) and (6.Q) are identical with (6.J), (6.G), (6.K) and (6.H) respectively. Thus they are exact. ☐

Proposition 6.4. *μ and ν are mutually inverse operations, i.e. we have $\nu_N(\mu_N(M))=M$ and $\mu_N(\nu_N(M))=M$, up to additive closure.*

Proof. This is an immediate consequence of 6.3. ☐

The following is standard in the theory of tilting mutation [RS91].

Lemma 6.5. *Let Λ be a ring, let Q be a projective Λ-module and consider an exact sequence $\Lambda\xrightarrow{f}Q'\xrightarrow{g}\operatorname{Cok}f\to 0$ where f is a left* $\operatorname{add}Q$*-approximation. If f is injective then $Q\oplus\operatorname{Cok}f$ is a tilting Λ-module of projective dimension ≤ 1.*

Proof. For the convenience of the reader we give a complete proof here. It is clear that $\operatorname{proj.dim}_\Lambda(Q\oplus\operatorname{Cok}f)\leq 1$ and it generates the derived category. We need only check that $\operatorname{Ext}^1_\Lambda(Q\oplus\operatorname{Cok}f,Q\oplus\operatorname{Cok}f)=0$. Applying $\operatorname{Hom}_\Lambda(-,Q)$, we have an exact sequence

$$\operatorname{Hom}_\Lambda(Q',Q)\xrightarrow{f\cdot}\operatorname{Hom}_\Lambda(\Lambda,Q)\to\operatorname{Ext}^1_\Lambda(\operatorname{Cok}f,Q)\to 0.$$

Since $(f\cdot)$ is surjective, we have $\operatorname{Ext}^1_\Lambda(\operatorname{Cok} f, Q) = 0$. Applying $\operatorname{Hom}_\Lambda(-, \operatorname{Cok} f)$, we have an exact sequence

$$\operatorname{Hom}_\Lambda(Q', \operatorname{Cok} f) \xrightarrow{f\cdot} \operatorname{Hom}_\Lambda(\Lambda, \operatorname{Cok} f) \to \operatorname{Ext}^1_\Lambda(\operatorname{Cok} f, \operatorname{Cok} f) \to 0.$$

Here $(f\cdot)$ is surjective since $\operatorname{Hom}_\Lambda(Q', Q') \xrightarrow{\cdot g} \operatorname{Hom}_\Lambda(\Lambda, Q')$ and $\operatorname{Hom}_\Lambda(\Lambda, Q') \xrightarrow{\cdot f} \operatorname{Hom}_\Lambda(\Lambda, \operatorname{Cok} f)$ are surjective. Thus we have $\operatorname{Ext}^1_\Lambda(\operatorname{Cok} f, \operatorname{Cok} f) = 0$. Consequently we have $\operatorname{Ext}^1_\Lambda(Q \oplus \operatorname{Cok} f, Q \oplus \operatorname{Cok} f) = 0$ since Q is projective. ☐

Recall from the introduction that given $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$ we define $[N]$ to be the two-sided ideal of $\operatorname{End}_R(M)$ consisting of morphisms $M \to M$ which factor through a member of $\operatorname{add} N$.

Theorem 6.6. *Let R be normal 3-CY$^-$ with modifying module M. Suppose $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$, denote $\Lambda = \operatorname{End}_R(M)$ and define $\Lambda_N := \Lambda/[N]$. If Λ_N is artinian then*
(1) *$T_1 := \operatorname{Hom}_R(M^*, \mu_N(M)^*)$ is a tilting Λ^{op}-module such that $\operatorname{End}_{\Lambda^{\mathrm{op}}}(T_1) \cong \operatorname{End}_R(\mu_N(M))^{\mathrm{op}}$. Thus $\operatorname{End}_R(M)^{\mathrm{op}}$ and $\operatorname{End}_R(\mu_N(M))^{\mathrm{op}}$ are derived equivalent, so $\operatorname{End}_R(M)$ and $\operatorname{End}_R(\mu_N(M))$ are derived equivalent.*
(2) *$T_2 := \operatorname{Hom}_R(M, \nu_N(M))$ is a tilting Λ-module such that $\operatorname{End}_\Lambda(T_2) \cong \operatorname{End}_R(\nu_N(M))$. Thus $\operatorname{End}_R(M)$ and $\operatorname{End}_R(\nu_N(M))$ are derived equivalent.*

Proof. (1) Denote $\mathbb{G} := \operatorname{Hom}_R(N^*, -)$ and $\Gamma := \operatorname{End}_R(N^*)$. Applying $\operatorname{Hom}_R(M^*, -)$ to (6.L) and $\operatorname{Hom}_\Gamma(\mathbb{G}M^*, -)$ to (6.N) gives a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & {}_\Gamma(\mathbb{G}M^*, \mathbb{G}M^*) \arrow[r] \arrow[d, equal] & {}_\Gamma(\mathbb{G}M^*, \mathbb{G}N_0^*) \arrow[r] \arrow[d, equal] & {}_\Gamma(\mathbb{G}M^*, \mathbb{G}K_0^*) \arrow[r] \arrow[d, equal] & \operatorname{Ext}^1_\Gamma(\mathbb{G}M^*, \mathbb{G}M^*) & \\
0 \arrow[r] & \operatorname{Hom}_R(M^*, M^*) \arrow[r, "\cdot a^*"] & \operatorname{Hom}_R(M^*, N_0^*) \arrow[r, "\cdot c^*"] & \operatorname{Hom}_R(M^*, K_0^*) \arrow[r] & \operatorname{Cok} \arrow[r] & 0
\end{tikzcd}$$

hence $\operatorname{Cok} \subseteq \operatorname{Ext}^1_\Gamma(\mathbb{G}M^*, \mathbb{G}M^*)$. We first claim that $\operatorname{Cok} = 0$. On one hand Γ is 3-CY$^-$ by 2.18 and so by 2.6 fl $\operatorname{Ext}^1_\Gamma(\mathbb{G}M^*, \mathbb{G}M^*) = 0$. On the other hand $\operatorname{Hom}_R(N^*, -)$ applied to (6.L) is exact (by 6.3) so Cok is a Λ_N^{op}-module. Since Λ_N^{op} is artinian so too is Cok, thus it has finite length. Hence $\operatorname{Cok} = 0$ and so

$$0 \to \operatorname{Hom}_R(M^*, M^*) \to \operatorname{Hom}_R(M^*, N_0^*) \to \operatorname{Hom}_R(M^*, K_0^*) \to 0$$

is exact. We now claim that this is a left-$\operatorname{add} Q$ approximation where $Q = \operatorname{Hom}_R(M^*, N^*)$. Simply applying $\operatorname{Hom}_{\Lambda^{\mathrm{op}}}(-, {}_R(M^*, N^*))$ to the above we obtain

$$\begin{tikzcd}
{}_{\Lambda^{\mathrm{op}}}({}_R(M^*, N_0^*), {}_R(M^*, N^*)) \arrow[r] \arrow[d, equal] & {}_{\Lambda^{\mathrm{op}}}({}_R(M^*, M^*), {}_R(M^*, N^*)) \arrow[d, equal] & \\
\operatorname{Hom}_R(N_0^*, N^*) \arrow[r] & \operatorname{Hom}_R(M^*, N^*) \arrow[r] & 0
\end{tikzcd}$$

where the bottom is just (6.O) and so is exact. Hence the top is exact and so by 6.5 $\operatorname{Hom}_R(M^*, N^*) \oplus \operatorname{Hom}_R(M^*, K_0^*) = \operatorname{Hom}_R(M^*, \mu_N(M)^*)$ is a tilting Λ^{op}-module. That $\operatorname{End}_{\Lambda^{\mathrm{op}}}(T_1) \cong \operatorname{End}_R(\mu_N(M)^*) = \operatorname{End}_R(\mu_N(M))^{\mathrm{op}}$ is just 2.5. Hence Λ^{op} and $\operatorname{End}_R(\mu_N(M))^{\mathrm{op}}$ are derived equivalent; it is well-known that this implies that Λ and $\operatorname{End}_R(\mu_N(M))$ are derived equivalent [R89, 9.1].
(2) Similarly to the above one can show that applying $\operatorname{Hom}_R(M, -)$ to (6.M) gives an exact sequence

$$0 \to \operatorname{Hom}_R(M, M) \to \operatorname{Hom}_R(M, N_1) \to \operatorname{Hom}_R(M, K_1^*) \to 0$$

which is an approximation. Hence by 6.5 $\operatorname{Hom}_R(M, N) \oplus \operatorname{Hom}_R(M, K_1^*) = \operatorname{Hom}_R(M, \nu(M))$ is a tilting module. The statement on endomorphism rings again holds by 2.5. ☐

Remark 6.7. Note that if R is finitely generated over a field k then Λ_N is artinian if and only if $\dim_k \Lambda_N < \infty$. Thus if the reader is willing to work over $\mathbb{C}$, they may replace the condition Λ_N is artinian by $\dim_{\mathbb{C}} \Lambda_N < \infty$ throughout.

Theorem 6.8. *Let R be normal 3-CY$^-$ with modifying module M. Suppose $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$, denote $\Lambda = \operatorname{End}_R(M)$ and define $\Lambda_N := \Lambda/[N]$. If Λ_N is artinian then*
(1) *If $\operatorname{add} N \neq \operatorname{add} M$ then $\operatorname{add} \mu_N(M) \neq \operatorname{add} M$.*
(2) *$\mu_N(M)$ is also a modifying module. Further*
- (a) *If M is maximal modifying then $\mu_N(M)$ is maximal modifying.*

  (b) *If* $\operatorname{End}_R(M)$ *is a NCCR then* $\operatorname{End}_R(\mu_N(M))$ *is a NCCR.*
  (c) *If* M *is a CT module and* N *is a generator then* $\mu_N(M)$ *is a CT module.*

$(1)'$ *If* $\operatorname{add} N \neq \operatorname{add} M$ *then* $\operatorname{add} \nu_N(M) \neq \operatorname{add} M$.
$(2)'$ $\nu_N(M)$ *is also a modifying module. Further*

  (a) *If* M *is maximal modifying then* $\nu_N(M)$ *is maximal modifying.*
  (b) *If* $\operatorname{End}_R(M)$ *is a NCCR then* $\operatorname{End}_R(\nu_N(M))$ *is a NCCR.*
  (c) *If* M *is a CT module and* N *is a generator then* $\nu_N(M)$ *is a CT module.*

Proof. (1) By 6.6(1)

$$0 \to \operatorname{Hom}_R(M^*, M^*) \to \operatorname{Hom}_R(M^*, N_0^*) \to \operatorname{Hom}_R(M^*, K_0^*) \to 0$$

is exact. If this splits then by reflexive equivalence M^* is a summand of N_0^*, contradicting $\operatorname{add} N \neq \operatorname{add} M$. Thus the above cannot split so $\operatorname{Hom}_R(M^*, K_0^*)$ cannot be projective, hence certainly $K_0^* \notin \operatorname{add} M^*$ and so $K_0 \notin \operatorname{add} M$. This implies $\operatorname{add} \mu_{M_1}(M) \neq \operatorname{add} M$.
(2) Using the derived equivalence in 6.6(1) together with 2.18 twice we deduce that $\operatorname{End}_R(\mu_N(M)) \in \operatorname{CM} R$, i.e. $\mu_N(M)$ is a modifying module. Alternatively just use 6.2.
(2a) follows from 4.14.
(2b) follows from (2a) and 5.14(1).
(2c) follows from (2b) and 5.14(2).
$(1)'$ Since the exact sequence in the proof of 6.6(2) cannot split, $K_1^* \notin \operatorname{add} M$.
$(2)'$ Is identical to (2). □

Remark 6.9. It is natural to ask under what circumstances the hypothesis Λ_N is artinian in 6.6, 6.8 holds. In the situation of 5.11 the answer seems to be related to the contractibility of the corresponding curves; we will come back to this question in future work.

One case where Λ_N is always artinian is when R has isolated singularities:

Lemma 6.10. *Suppose* R *is normal 3-CY*$^-$. *Let* M *be a modifying module with* $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$, *denote* $\Lambda = \operatorname{End}_R(M)$ *and set* $\Lambda_N = \Lambda/[N]$. *Then*
(1) $\dim_R \Lambda_N \leq 1$.
(2) $\operatorname{depth}_{R_\mathfrak{m}} (\Lambda_N)_\mathfrak{m} \leq 1$ *for all* $\mathfrak{m} \in \operatorname{Max} R$.
(3) *If* R *is an isolated singularity then* Λ_N *is artinian.*
(4) *If* $\operatorname{proj.dim}_\Lambda \Lambda_N < \infty$ *then* $\operatorname{inj.dim}_{\Lambda_N} \Lambda_N \leq 1$.

Proof. (1) We know that $\dim \operatorname{Sing} R \leq 1$ since R is normal. Now because $N \in \operatorname{add} M$ we may write $\operatorname{End}_R(M^n) = \operatorname{End}_R(N \oplus T)$ say. Then since $\operatorname{End}_R(M)$ is Morita equivalent to $\operatorname{End}_R(M^n)$, we have that Λ_N is Morita equivalent to $\operatorname{End}_R(N \oplus T)/[N]$ where $[N]$ again denotes the two-sided ideal of those morphisms which factor through a member of $\operatorname{add} N$. Let $\mathfrak{p} \in \operatorname{Spec} R$ with $\dim R/\mathfrak{p} > 1$. We show that $(\operatorname{End}_R(N \oplus T)/[N])_\mathfrak{p} = 0$ as via the Morita equivalence this shows that $(\Lambda_N)_\mathfrak{p} = 0$ and hence $\dim_R \Lambda_N \leq 1$. But $(\operatorname{End}_R(N \oplus T)/[N])_\mathfrak{p} = \operatorname{End}_{R_\mathfrak{p}}(N_\mathfrak{p} \oplus T_\mathfrak{p})/[N]_\mathfrak{p}$ and $\operatorname{End}_{R_\mathfrak{p}}(N_\mathfrak{p} \oplus T_\mathfrak{p})$ is some matrix ring over $R_\mathfrak{p}$ since $N_\mathfrak{p} \oplus T_\mathfrak{p}$ is a reflexive module over a regular local ring $R_\mathfrak{p}$ with $\dim R_\mathfrak{p} \leq 1$, thus is a free $R_\mathfrak{p}$-module. Obviously $[N] \subseteq \operatorname{End}_R(N \oplus T)$ contains the idempotent $e : N \oplus T \xrightarrow{\left(\begin{smallmatrix}1&0\\0&0\end{smallmatrix}\right)} N \oplus T$. Now $e_\mathfrak{p} : N_\mathfrak{p} \oplus T_\mathfrak{p} \xrightarrow{\left(\begin{smallmatrix}1&0\\0&0\end{smallmatrix}\right)} N_\mathfrak{p} \oplus T_\mathfrak{p}$ is non-zero since $N_\mathfrak{p} \neq 0$, thus $[N]_\mathfrak{p}$ contains the non-zero idempotent $e_\mathfrak{p}$. But for matrix algebras every non-zero idempotent generates the ring. Consequently $(\Lambda_N)_\mathfrak{p} = 0$.
(2) is immediate from (1).
(3) If R is isolated then by (1) $\dim_R \Lambda_N = 0$ and so Λ_N is supported only at a finite number of maximal ideals. Hence Λ_N has finite length and so Λ_N is artinian.
(4) Notice that Λ is 3-CY$^-$ by 2.18. Hence the assertion follows from [IR08, 5.5(3)] for 3-CY algebras, which is also valid for 3-CY$^-$ algebras under the assumption that $\operatorname{proj.dim}_\Lambda \Lambda_N < \infty$. □

We now show that mutation does not change the factor algebra Λ_N. Suppose M is modifying and N is such that $0 \neq \operatorname{add} N \subseteq \operatorname{add} M$, and consider an exchange sequence

$$0 \to K_0 \xrightarrow{c} N_0 \xrightarrow{a} M.$$

We know by definition that a is a right $\operatorname{add} N$-approximation, and by (6.J) that c is a left $\operatorname{add} N$-approximation.

Since Λ_N is by definition $\operatorname{End}_R(M)$ factored out by the ideal of all morphisms $M \to M$ which factor through a module in $\operatorname{add} N$, in light of the universal property of the map a, this ideal is the just the ideal I_a of all morphisms $M \to M$ which factor as xa where x is some morphism $M \to N_0$. Thus $\Lambda_N = \operatorname{End}_R(M)/I_a$.

On the other hand taking the choice $\mu_N(M) = K_0 \oplus N$ coming from the above exchange sequence, Λ'_N is by definition $\operatorname{End}_R(\mu_N(M)) = \operatorname{End}_R(K_0 \oplus N)$ factored out by the ideal of all morphisms $K_0 \oplus N \to K_0 \oplus N$ which factor through a module in $\operatorname{add} N$. Clearly this is just $\operatorname{End}_R(K_0)$ factored out by those morphisms which factor through $\operatorname{add} N$. In light of the universal property of the map c, $\Lambda'_N = \operatorname{End}_R(K_0)/I_c$ where I_c is the ideal of all morphisms $K_0 \to K_0$ which factor as cy where y is some morphism $K_0 \to N_0$.

Theorem 6.11. *With the choice of exchange sequence as above, we have $\Lambda_N \cong \Lambda'_N$. In particular*
(1) Λ'_N is independent of the choice of exchange sequence, up to isomorphism.
(2) Λ_N is artinian if and only if Λ'_N is artinian.

Proof. We construct a map $\Lambda_N = \operatorname{End}_R(M)/I_a \to \operatorname{End}_R(K_0)/I_c = \Lambda'_N$ as follows: given $f \in \operatorname{End}_R(M)$ we have

$$\begin{tikzcd}
0 \arrow[r] & K_0 \arrow[r, "c"] \arrow[d, dotted, "\exists\, h_f"] & N_0 \arrow[r, "a"] \arrow[d, dotted, "\exists\, g_f"] & M \arrow[d, "f"] \\
0 \arrow[r] & K_0 \arrow[r, "c"] & N_0 \arrow[r, "a"] & M
\end{tikzcd}$$

where the g_f exists (non-uniquely) since a is an approximation. Define $\alpha : \Lambda_J \to \Lambda'_J$ by $\alpha(f + I_a) = h_f + I_c$. Note first that this is independent of the choice of g_f, for if g'_f is another morphism such that $g'_f a = af$ then

$$\begin{tikzcd}
0 \arrow[r] & K_0 \arrow[r, "c"] \arrow[d, dotted, "\exists\, h'_f"] & N_0 \arrow[r, "a"] \arrow[d, "g'_f"] & M \arrow[d, "f"] \\
0 \arrow[r] & K_0 \arrow[r, "c"] & N_0 \arrow[r, "a"] & M
\end{tikzcd}$$

From this $(g_f - g'_f)a = a(f-f) = 0$ and so there exists $y : N_0 \to K_0$ such that $yc = g_f - g'_f$. Hence $cyc = c(g_f - g'_f) = (h_f - h'_f)c$ and so since c is a monomorphism $cy = h_f - h'_f$. Consequently $h_f + I_c = h'_f + I_c$ and so α is independent of the choice of g_f.

Due to the independence of the choice of g_f it follows immediately that α is an algebra homomorphism provided that it is well-defined. But α is well defined since if $f_1 + I_a = f_2 + I_a$ then there exists $x : M \to N_0$ such that $xa = f_1 - f_2$. Thus $(ax - g_{f_1} + g_{f_2})a = 0$ so there exists $y : N_0 \to K_0$ such that $yc = ax - g_{f_1} + g_{f_2}$. This implies $-cyc = c(g_{f_1} - g_{f_2}) = (h_{f_1} - h_{f_2})c$ so since c is a monomorphism $-cy = h_{f_1} - h_{f_2}$ and so $\alpha(f_1 + I_a) = h_{f_1} + I_c = h_{f_2} + I_c = \alpha(f_2 + I_a)$ as required.

We now show that α is onto. Let $t : K_0 \to K_0$ be any morphism then on dualizing we have

$$\begin{tikzcd}
0 \arrow[r] & M^* \arrow[r, "a^*"] & N_0^* \arrow[r, "c^*"] & K_0^* \\
0 \arrow[r] & M^* \arrow[r, "a^*"] \arrow[u, dotted, "\exists\, r"] & N_0^* \arrow[r, "c^*"] \arrow[u, dotted, "\exists\, s"] & K_0^* \arrow[u, "t^*"]
\end{tikzcd}$$

where the rows are exact by (6.L), s exists (non-uniquely) by (6.N) and r exists since a^* is the kernel of c^*. Dualizing back gives a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & K_0 \arrow[r, "c"] \arrow[d, "t"] & N_0 \arrow[r, "a"] \arrow[d, dotted, "s^*"] & M \arrow[d, dotted, "r^*"] \\
0 \arrow[r] & K_0 \arrow[r, "c"] & N_0 \arrow[r, "a"] & M
\end{tikzcd}$$

and so it follows that $\alpha(r^* + I_a) = t + I_c$.

Finally, α is injective. To see this suppose $\alpha(f_1+I_a)=\alpha(f_2+I_a)$ then there exists $y:N_0\to K_0$ such that $h_{f_1}-h_{f_2}=cy$. Hence $c(yc-g_{f_1}+g_{f_2})=0$ and so since by (6.J)

$$0\to \mathrm{Hom}_R(M,N_0)\xrightarrow{a\cdot}\mathrm{Hom}_R(N_0,N_0)\xrightarrow{c\cdot}\mathrm{Hom}_R(K_0,N_0)\to 0.$$

is exact there exists $x:M\to N_0$ such that $ax=yc-g_{f_1}+g_{f_2}$. This implies that $a(xa-f_1+f_2)=0$, but denoting $\mathbb{F}=\mathrm{Hom}_R(N,-)$ we have a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \mathrm{Hom}(\mathbb{F}M,\mathbb{F}M) \arrow[r] \arrow[d, equal] & \mathrm{Hom}(\mathbb{F}N_0,\mathbb{F}M) \arrow[r] \arrow[d, equal] & \mathrm{Hom}(\mathbb{F}K_0,\mathbb{F}M) \arrow[d, equal] \\
 & \mathrm{Hom}_R(M,M) \arrow[r, "a\cdot"] & \mathrm{Hom}_R(N_0,M) \arrow[r, "c\cdot"] & \mathrm{Hom}_R(K_0,M)
\end{tikzcd}$$

from which we see that $a\cdot$ is injective. Consequently $xa-f_1+f_2=0$ and so $f_1+I_a=f_2+I_a$, as required. ☐

6.2. **Complete Local Case.** From now on we assume that R is a *complete local* normal Gorenstein three-dimensional ring, then since we have Krull–Schmidt decompositions we can say more than in the previous section. Note that with these assumptions R is automatically 3-CY$^-$ by 2.16. For a modifying module M we write

$$M=M_1\oplus\ldots\oplus M_n=\bigoplus_{i\in I}M_i$$

as its Krull–Schmidt decomposition into indecomposable submodules, where $I=\{1,\ldots,n\}$. With the new assumption on R we may take minimal approximations and so the setup in the previous section can be simplified: for $\emptyset\neq J\subseteq I$ set $M_J:=\oplus_{j\in J}M_j$ and $\frac{M}{M_J}:=\oplus_{i\in I\backslash J}M_i$. Then

(a) Denote $L_0\xrightarrow{a}M_J$ to be a right $\mathrm{add}(\frac{M}{M_J})$-approximation of M_J which is *right minimal*. If $\frac{M}{M_J}$ contains R as a summand then necessarily a is surjective.

(b) Similarly denote $L_1^*\xrightarrow{b}M_J^*$ to be a right $\mathrm{add}(\frac{M^*}{M_J^*})$-approximation of M_J^* which is *right minimal*. Again if $\frac{M}{M_J}$ contains R as a summand then b is injective.

Recall that a morphism $a:X\to Y$ is called *right minimal* if any $f\in\mathrm{End}_R(X)$ satisfying $a=fa$ is an automorphism. In what follows we denote the kernels of the above right minimal approximations by

$$0\to C_0\xrightarrow{c}L_0\xrightarrow{a}M_J\quad\text{and}\quad 0\to C_1\xrightarrow{d}L_1^*\xrightarrow{b}M_J^*.$$

This recovers the mutations from the previous subsection:

Lemma 6.12. *With notation as above,* $\mu_{\frac{M}{M_J}}(M)=\frac{M}{M_J}\oplus C_0$ *and* $\nu_{\frac{M}{M_J}}(M)=\frac{M}{M_J}\oplus C_1^*$.

Proof. There is an exact sequence

$$0\to C_0\xrightarrow{(c\ 0)}L_0\oplus\frac{M}{M_J}\xrightarrow{\left(\begin{smallmatrix}a&0\\0&1\end{smallmatrix}\right)}M_J\oplus\frac{M}{M_J}=M\to 0$$

with $L_0\oplus\frac{M}{M_J}\in\mathrm{add}\,\frac{M}{M_J}$ such that $\mathrm{Hom}_R(\frac{M}{M_J},-)$ is exact. The result follows from 6.1. The other statement is similar. ☐

Since we have minimal approximations from now on we *define* our mutations in terms of them. We thus define $\mu_J(M):=C_0\oplus\frac{M}{M_J}$ and $\nu_J(M):=C_1^*\oplus\frac{M}{M_J}$. When $J=\{i\}$ we often write ν_i and μ_i instead of $\nu_{\{i\}}$ and $\mu_{\{i\}}$ respectively.

We now characterize when the mutation of an indecomposable summand is trivial. Note that using the above definition of mutation involving minimal approximations, μ and ν are now inverse operations up to isomorphism, not just additive closure. This strengthens 6.4.

Theorem 6.13. *Suppose R is a complete local normal Gorenstein three-dimensional ring and let M be a MM module with indecomposable summand M_i. Denote $\Lambda=\mathrm{End}_R(M)$, let e_i be the idempotent corresponding to M_i and denote $\Lambda_i=\Lambda/\Lambda(1-e_i)\Lambda$. Suppose Λ_i is not artinian, then $\mu_i(M)=\nu_i(M)=M$.*

Proof. We first show that $\operatorname{proj.dim}_\Lambda \Lambda_i < \infty$. The sequence

$$0 \to C_0 \to L_0 \to M_i$$

gives

$$0 \to \operatorname{Hom}_R(M, C_0) \to \operatorname{Hom}_R(M, L_0) \to \operatorname{Hom}_R(M, M_i) \to \Lambda_i \to 0$$

where $\operatorname{Hom}_R(M, L_0)$ and $\operatorname{Hom}_R(M, M_i)$ are projective Λ-modules. But C_0 is a modifying module by 6.2 so by 4.6 we know that $\operatorname{proj.dim}_\Lambda \operatorname{Hom}_R(M, C_0) \leq 1$ and hence certainly $\operatorname{proj.dim}_\Lambda \Lambda_i \leq 3$.

Now it is always true that $\operatorname{depth} \Lambda_i \leq \dim \Lambda_i \leq \operatorname{inj.dim}_{\Lambda_i} \Lambda_i$ by [GN02, 3.5] (see [IR08, 2.1]). Since $\operatorname{proj.dim}_\Lambda \Lambda_i \leq 3$ by 6.10(4) we know $\operatorname{inj.dim}_{\Lambda_i} \Lambda_i \leq 1$. This being the case $\operatorname{depth} \Lambda_i = \operatorname{inj.dim}_{\Lambda_i} \Lambda_i$ by Ramras [Ram, 2.15]. If $\dim \Lambda_i = 0$ then $\operatorname{Supp}_R \Lambda_i$ contains only the maximal ideal and so Λ_i has finite length, contradicting the assumption that Λ_i is not artinian. Thus $\operatorname{depth} \Lambda_i = \dim \Lambda_i = \operatorname{inj.dim}_{\Lambda_i} \Lambda_i = 1$.

Now Λ is a Gorenstein R-order by 2.17 and 2.18 so by Auslander–Buchsbaum 2.13, since $\operatorname{proj.dim}_\Lambda \Lambda_i < \infty$ necessarily $\operatorname{proj.dim}_\Lambda \Lambda_i = 3 - \operatorname{depth} \Lambda_i = 2$. Thus we have a minimal projective resolution

$$0 \to P_2 \to P_1 \xrightarrow{f} \Lambda e_i \to \Lambda_i \to 0. \tag{6.R}$$

where f is a minimal right $\operatorname{add} \Lambda(1-e_i)$-approximation since it is a projective cover of $\Lambda(1-e_i)\Lambda e_i$. By [IR08, 3.4(5)] we have

$$\operatorname{Ext}^2_\Lambda(\Lambda_i, \Lambda) \cong \operatorname{Ext}^2_R(\Lambda_i, R)$$

and this is a projective Λ_i^{op}-module by [GN02, 1.1(3)]. Since $\operatorname{Ext}^2_R(\operatorname{Ext}^2_R(\Lambda_i, R), R) \cong \Lambda_i$ and Λ_i is a local ring, we have $\operatorname{Ext}^2_R(\Lambda_i, R) \cong \Lambda_i$ as Λ_i^{op}-modules. Applying $\operatorname{Hom}_\Lambda(-, \Lambda)$ to (6.R) gives an exact sequence

$$\operatorname{Hom}_\Lambda(P_1, \Lambda) \to \operatorname{Hom}_\Lambda(P_2, \Lambda) \to \Lambda_i \to 0$$

which gives a minimal projective presentation of the Λ^{op}-module Λ_i. Thus we have $\operatorname{Hom}_\Lambda(P_2, \Lambda) \cong e_i\Lambda$ and $P_2 \cong \Lambda e_i$.

Under the equivalence $\operatorname{Hom}_R(M, -) : \operatorname{add} M \to \operatorname{proj} \Lambda$, the sequence (6.R) corresponds to a complex

$$0 \to M_i \xrightarrow{h} L_0 \xrightarrow{g} M_i$$

with g a minimal right $\operatorname{add} \frac{M}{M_i}$-approximation. Since the induced morphism $M_i \to \operatorname{Ker} g$ is sent to an isomorphism under the reflexive equivalence $\operatorname{Hom}_R(M, -) : \operatorname{ref} R \to \operatorname{ref} \Lambda$ (2.5(4)), it is an isomorphism and so $h = \ker g$. Consequently we have $\mu_i(M) = \frac{M}{M_i} \oplus M_i = M$. This implies that $\nu_i(M) = M$ by 6.4. $\square$

Summarizing what we have proved, in the complete local case we have

Theorem 6.14. *Suppose R is a complete local normal three-dimensional Gorenstein ring with maximal modifying module M. Denote $\Lambda = \operatorname{End}_R(M)$, let M_i be an indecomposable summand of M and consider $\Lambda_i := \Lambda/\Lambda(1-e_i)\Lambda$ where e_i is the idempotent in Λ corresponding to M_i. Then*
(1) *If Λ_i is not artinian then $\mu_i(M) = \nu_i(M) = M$.*
(2) *If Λ_i is artinian then $\mu_i(M) = \nu_i(M)$ and this is not equal to M.*
In either case denote $\mu_i := \mu_i = \nu_i$ then it is also true that
(3) *$\mu_i\mu_i(M) = M$.*
(4) *$\mu_i(M)$ is a maximal modifying module.*
(5) *$\operatorname{End}_R(M)$ and $\operatorname{End}_R(\mu_i(M))$ are derived equivalent, via the tilting $\operatorname{End}_R(M)$-module $\operatorname{Hom}_R(M, \mu_i(M))$.*

Proof. (1) is 6.13.
(2) M, $\mu_i(M)$ and $\nu_i(M)$ are all MM modules by 6.8, thus by 4.15(1) both $\operatorname{Hom}_R(M, \mu_i(M))$ and $\operatorname{Hom}_R(M, \nu_i(M))$ are tilting $\operatorname{End}_R(M)$ modules. But since $\mu_i(M) \neq M$ and $\nu_i(M) \neq M$ (by 6.8) neither of these tilting modules equal $\operatorname{Hom}_R(M, M)$. Further by construction $\operatorname{Hom}_R(M, \mu_i(M))$ and $\operatorname{Hom}_R(M, \nu_i(M))$ share all summands except possibly one, thus by Riedtmann-Schofield [RS91] necessarily it is true that $\operatorname{Hom}_R(M, \mu_i(M)) \cong \operatorname{Hom}_R(M, \nu_i(M))$. By reflexive equivalence we deduce that $\mu_i(M) \cong \nu_i(M)$.

The remainder is trivially true in the case when Λ_i is not artinian (by 6.13), thus we may assume that Λ_i is artinian. Now $\mu_i\mu_i(M) = \mu_i(\nu_i M) = \mu_i(C_1^* \oplus \frac{M}{M_i}) = K \oplus \frac{M}{M_i}$ where K is the kernel of a minimal left add $\frac{M}{M_i}$-approximation of C_1^*. But by a similar version of (6.M) and (6.P) we know $K = M_i$. Thus (3) follows. (4) is contained in 6.8 and (5) is 6.6(2). □

Example 6.15. Consider the group $\frac{1}{2}(1,1,0) \oplus \frac{1}{2}(0,1,1)$ and let $R = k[[x,y,z]]^G$. We know by 5.8 that $M = k[[x,y,z]]$ is a CT R-module, and in this example it decomposes into 4 summands $R \oplus M_1 \oplus M_2 \oplus M_3$ with respect to the characters of G. The quiver of $\mathrm{End}_R(M)$ is the McKay quiver

$$\begin{tikzcd}
M_1 \arrow[r, shift left, "y"] \arrow[d, shift left, "x"] \arrow[dr, shift left, "z"] & M_2 \arrow[l, shift left, "y"] \arrow[d, shift left, "x"] \arrow[dl, shift left, "z"] \\
R \arrow[u, shift left, "x"] \arrow[r, shift left, "y"] \arrow[ur, shift left, "z"] & M_3 \arrow[l, shift left, "y"] \arrow[u, shift left, "x"] \arrow[ul, shift left, "z"]
\end{tikzcd}$$

and so to mutate at M_2 it is clear that the relevant approximation is

$$R \oplus M_1 \oplus M_3 \xrightarrow{\begin{pmatrix} x \\ y \\ z \end{pmatrix}} M_2 \to 0$$

Thus the mutation at vertex M_2 changes $M = R \oplus M_1 \oplus M_2 \oplus M_3$ into $R \oplus M_1 \oplus K_2 \oplus M_3$ where K_2 is the kernel of the above map which (by counting ranks) has rank 2. On the level of quivers of the endomorphism rings, this induces the mutation



Due to the relations in the algebra $\mathrm{End}_R(\mu_2(M))$ (which we suppress), the mutation at M_1 and M_3 in the new quiver are trivial, thus in $\mathrm{End}_R(\mu_2(M))$ the only vertex we can mutate at is K_2, which gives us back our original M. By the symmetry of the situation we obtain the mutation graph

$$\begin{tikzcd}
 & R\oplus M_1\oplus M_2\oplus K_3 \arrow[d, dash, dotted, "\mu_3"] & \\
R\oplus K_1\oplus M_2\oplus M_3 \arrow[r, dash, dotted, "\mu_1"] & R\oplus M_1\oplus M_2\oplus M_3 \arrow[r, dash, dotted, "\mu_2"] & R\oplus M_1\oplus K_2\oplus M_3
\end{tikzcd}$$

We remark that mutating at any of the decomposable modules $M_1 \oplus M_2$, $M_1 \oplus M_3$, $M_2 \oplus M_3$ or $M_1 \oplus M_2 \oplus M_3$ gives a trivial mutation.

Acknowledgements. We thank Michel Van den Bergh for valuable suggestions. We thank Ryo Takahashi and Yuji Yoshino for stimulating discussions.

References

[Aus78] M. Auslander, *Functors and morphisms determined by objects.* Representation theory of algebras (Proc. Conf., Temple Univ., Philadelphia, Pa., 1976), pp. 1–244. Lecture Notes in Pure Appl. Math., Vol. 37, Dekker, New York, 1978.

[Aus84] ______, *Isolated singularities and existence of almost split sequences.* Representation theory, II (Ottawa, Ont., 1984), 194–242, Lecture Notes in Math., 1178, Springer, Berlin, 1986.

[Aus86] ______, *Rational singularities and almost split sequences.* Trans. Amer. Math. Soc. **293** (1986), no. 2, 511–531.

[AR] M. Auslander and I. Reiten, *Almost split sequences for Cohen-Macaulay modules*, Math. Ann. **277** (1987), 345–349.

[BBD] A. A. Beilinson, J. Bernstein and P. Deligne, *Faisceaux pervers*, Analysis and topology on singular spaces, I (Luminy, 1981), 5–171, Astérisque, **100**, Soc. Math. France, Paris, 1982.

[Bou68] N. Bourbaki, *Groupe et Algebre de Lie*, Chapter 5, Hermann, Paris, 1968.

[BKR] T. Bridgeland, A. King and M. Reid, *The McKay correspondence as an equivalence of derived categories*, J. Amer. Math. Soc. **14** (2001), no. 3, 535–554.

[BH] W. Bruns and J. Herzog, *Cohen-Macaulay rings. Rev. ed.*, Cambridge Studies in Advanced Mathematics. **39** xiv+453 pp.

[BIKR] I. Burban, O. Iyama, B. Keller and I. Reiten, *Cluster tilting for one-dimensional hypersurface singularities*, Adv. Math. **217** (2008), no. 6, 2443–2484.

[CB] W. Crawley-Boevey, *On the exceptional fibres of Kleinian singularities*, Amer. J. Math. **122** (2000), no. 5, 1027–1037.

[CE99] H. Cartan and S. Eilenberg, *Homological algebra.* Reprint of the 1956 original. Princeton Landmarks in Mathematics. Princeton University Press, 1999.

[C02] J-C. Chen, *Flops and equivalences of derived categories for threefolds with only terminal Gorenstein singularities*, J. Differential Geom. **61** (2002), no. 2, 227–261.

[CR90] C. W. Curtis and I. Reiner, *Methods of representation theory. Vol. I. With applications to finite groups and orders.* Reprint of the 1981 original. John Wiley & Sons, Inc., New York, 1990.

[Ei95] D. Eisenbud, *Commutative algebra. With a view toward algebraic geometry*, Graduate Texts in Mathematics, **150** Springer-Verlag, New York, 1995. xvi+785 pp.

[GLS] C. Geiss, B. Leclerc and J. Schröer, *Rigid modules over preprojective algebras*, Invent. Math. **165** (2006), no. 3, 589–632.

[GN02] S. Goto and K. Nishida, *Towards a theory of Bass numbers with application to Gorenstein algebras*, Colloq. Math. **91** (2002), no. 2, 191–253.

[Iya07] O. Iyama, *Higher-dimensional Auslander–Reiten theory on maximal orthogonal subcategories*, Adv. Math. **210** (2007), no. 1, 22–50.

[IR08] O. Iyama and I. Reiten, *Fomin-Zelevinsky mutation and tilting modules over Calabi-Yau algebras*, Amer. J. Math. **130** (2008), no. 4, 1087–1149.

[IT] O. Iyama and R. Takahashi, *Tilting and cluster tilting for quotient singularities*, in preparation.

[IW08] O. Iyama and M. Wemyss, *The classification of special Cohen Macaulay modules*, Math. Z. **265** (2010), no.1, 41–83.

[Lam] T. Y. Lam, *Lectures on modules and rings*, Graduate Texts in Mathematics, **189** Springer-Verlag, New York, 1999. xxiv+557 pp.

[Mat] H. Matsumura, *Commutative ring theory* (Second edition), Cambridge Studies in Advanced Mathematics, **8**, Cambridge University Press, Cambridge, 1989.

[M03] J. Miyachi, *Recollement and tilting complexes*, J. Pure Appl. Algebra **183** (2003) 245–273.

[O76] A. Ooishi, *Matlis duality and the width of a module*, Hiroshima Math. J. **6** (1976), 573–587.

[Ram] M. Ramras, *Maximal orders over regular local rings of dimension two*, Trans. Amer. Math. Soc. **142** (1969), 457–479.

[RV89] I. Reiten and M. Van den Bergh, *Two-dimensional tame and maximal orders of finite representation type.*, Mem. Amer. Math. Soc. **408** (1989), vii+72pp.

[R89] J. Rickard, *Morita theory for derived categories*, J. London Math. Soc. (2) **39** (1989), 436–456.

[RS91] C. Riedtmann and A. H. Schofield, *On a simplicial complex associated with tilting modules*, Comment. Math. Helv. **66** (1991), no. 1, 70–78.

[V04a] M. Van den Bergh, *Three-dimensional flops and noncommutative rings*, Duke Math. J. **122** (2004), no. 3, 423–455.

[V04b] ———, *Non-commutative crepant resolutions*, The legacy of Niels Henrik Abel, 749–770, Springer, Berlin, 2004.

[Yos90] Y. Yoshino, *Cohen-Macaulay modules over Cohen-Macaulay rings*, London Mathematical Society Lecture Note Series, 146. Cambridge University Press, Cambridge, 1990.

Osamu Iyama, Graduate School of Mathematics, Nagoya University, Chikusa-ku, Nagoya, 464-8602, Japan
E-mail address: `iyama@math.nagoya-u.ac.jp`

Michael Wemyss, School of Mathematics, James Clerk Maxwell Building, The King's Buildings, Mayfield Road, Edinburgh, EH9 3JZ, UK.
E-mail address: `wemyss.m@googlemail.com`